Draft version August 4, 2008
Preprint typeset using LaTeX style emulateapj v. 10/09/06

arXiv:0808.0471v1 [astro-ph] 4 Aug 2008

DISKS AROUND BROWN DWARFS IN THE σ ORIONIS CLUSTER[1]

K. L. Luhman[2], J. Hernández[3,4], J. J. Downes[4,5], L. Hartmann[3], and C. Briceño[4]

Draft version August 4, 2008

ABSTRACT

We have performed a census of circumstellar disks around brown dwarfs in the σ Ori cluster using all available images from the Infrared Array Camera onboard the *Spitzer Space Telescope.* To search for new low-mass cluster members with disks, we have measured photometry for all sources in the *Spitzer* images and have identified the ones that have red colors that are indicative of disks. We present five promising candidates, which may consist of two brown dwarfs, two stars with edge-on disks, and a low-mass protostar if they are bona fide members. Spectroscopy is needed to verify the nature of these sources. We have also used the *Spitzer* data to determine which of the previously known probable members of σ Ori are likely to have disks. By doing so, we measure disk fractions of $\sim 40\%$ and $\sim 60\%$ for low-mass stars and brown dwarfs, respectively. These results are similar to previous estimates of disk fractions in IC 348 and Chamaeleon I, which have roughly the same median ages as σ Ori ($\tau \sim 3$ Myr). Finally, we note that our photometric measurements and the sources that we identify as having disks differ significantly from those of other recent studies that analyzed the same *Spitzer* images. For instance, previous work has suggested that the T dwarf S Ori 70 is redder than typical field dwarfs, which has been cited as possible evidence of youth and cluster membership. However, we find that this object is only slightly redder than the reddest field dwarfs in $[3.6]-[4.5]$ (1.56 ± 0.07 vs. 0.93–1.46). We measure a larger excess in $[3.6]-[5.8]$ (1.75 ± 0.21 vs. 0.87–1.19), but the flux at 5.8 μm may be overestimated because of the low signal-to-noise ratio of the detection. Thus, the *Spitzer* data do not offer strong evidence of youth and membership for this object, which is the faintest and coolest candidate member of σ Ori that has been identified to date.

Subject headings: accretion disks — planetary systems: protoplanetary disks — stars: formation — stars: low-mass, brown dwarfs — stars: pre-main sequence

1. INTRODUCTION

The lifetime of an accretion disk around a young star represents a fundamental constraint on the amount of time available for the formation of giant planets. The typical lifetimes of disks are estimated by comparing the prevalence of disks among clusters that span a range of ages ($\tau \sim$ 1–10 Myr). By measuring disk fractions as a function of stellar mass and star-forming environment as well as age, the influence of these two factors on disk lifetimes can be characterized.

Disk fractions are usually measured through infrared (IR) photometry of young clusters ($\lambda > 3$ μm) and the identification of the stars that exhibit excess emission from cool dust. Observations of this kind were first performed with the *Infrared Astronomical Satellite* and ground-based near-IR cameras (Kenyon & Hartmann 1995; Haisch, Lada, & Lada 2001) and have made rapid progress in recent years with the *Spitzer Space Telescope* (Werner et al. 2004). Because *Spitzer* is exceptionally well-suited for detecting disks around members of young clusters (Allen et al. 2004; Gutermuth et al. 2004; Megeath et al. 2004; Muzerolle et al. 2004), it has been used extensively for measuring the disk fractions of stars (Megeath et al. 2005; Lada et al. 2006; Carpenter et al. 2006; Sicilia-Aguilar et al. 2006; Hernández et al. 2007a,b, 2008; Muench et al. 2007; Dahm & Hillenbrand 2007; Barrado y Navascués et al. 2007; Damjanov et al. 2007; Luhman et al. 2008). *Spitzer*'s unique sensitivity to faint IR sources has enabled measurements of disk fractions well into the substellar regime in nearby star-forming regions (d = 150–300 pc, Luhman et al. 2005, 2006, 2008; Guieu et al. 2007).

The cluster of young stars associated with the O star σ Ori A has been thoroughly searched for low-mass stars and brown dwarfs (e.g., Béjar et al. 2001; Barrado y Navascués et al. 2002; Martín et al. 2001; Zapatero Osorio et al. 2002a,b). As a result, it is an attractive target for measuring the disk fraction among low-mass objects. Several recent studies have sought to do this with *Spitzer*. Hernández et al. (2007a) performed *Spitzer* imaging of most of the σ Ori cluster and measured the disk fraction as a function of mass down to ~ 0.1 $M_\odot$ for a sample of ~ 300 probable members (see also Caballero (2006)). Caballero et al. (2007) identified a sample of brown dwarf candidates from optical and near-IR data and used the *Spitzer* images from Hernández et al. (2007a) to estimate a disk fraction for those sources, arriving at a value of $\sim$ 50%. Through further analysis of those *Spitzer* data, Zapatero Osorio et al. (2007) found that 6 of 12 brown dwarf candidates exhibited excess emission at 8 μm. Scholz & Jayawardhana (2008) obtained deeper *Spitzer* images of 18 of the faintest candidate members and reported a disk fraction of 29%. In addition to measuring disk fractions, the *Spitzer* data have been used to assess the youth and membership of the coolest candidate member of the cluster, the T dwarf S Ori 70 (Zapatero Osorio et al. 2002a). Zapatero Osorio et al. (2008) found that it is redder than typical field dwarfs in $[3.6] - [4.5]$ in the images from Hernández et al. (2007a). Similarly, Scholz & Jayawardhana (2008) suggested that S Ori 70 could be anomalously red in $[3.6] - [4.5]$ and $[3.6] - [5.8]$ based on their deeper images, which they attributed to a disk or low surface gravity. In either case, the apparent color excesses would comprise evidence of youth for this object, whose membership in the σ Ori cluster has been questioned (Martín & Zapatero Osorio 2003; Burgasser et al. 2004).

---

[1] Based on observations performed with the *Spitzer Space Telescope*.

[2] Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802; kluhman@astro.psu.edu.

[3] Department of Astronomy, The University of Michigan, Ann Arbor, MI 48109.

[4] Centro de Investigaciones de Astronomía, Apartado Postal. 264, Mérida 5101-A, Venezuela.

[5] Escuela de Física, Universidad Central de Venezuela, Apartado Postal 47586, Caracas 1041-A, Venezuela

The substellar population of the σ Ori cluster extends down to and below the detection limits of the images that have been obtained by *Spitzer*. Thus, determining whether these objects have disks is a challenging task. For instance, Zapatero Osorio et al. (2007) and Scholz & Jayawardhana (2008) disagree on whether the *Spitzer* data show evidence of disks for half of the sources considered by both studies. In this paper, we seek to accurately characterize the disk population among low-mass stars and brown dwarfs in σ Ori through an analysis of all *Spitzer* images of this cluster that includes careful treatment of errors and biases. We begin by summarizing the *Spitzer* observations and our data reduction methods (§ 2). We then use these data to search for new low-mass members of σ Ori that have disks (§ 3) and to estimate the disk fraction for the substellar members of the cluster (§ 4).

2. OBSERVATIONS

For our study of disks around brown dwarfs in the σ Ori cluster, we use images at 3.6, 4.5, 5.8, and 8.0 μm obtained with *Spitzer*'s Infrared Array Camera (IRAC; Fazio et al. 2004). We consider all IRAC observations that have been performed in this region, which consist of images analyzed by Hernández et al. (2007a) and Scholz & Jayawardhana (2008). Henceforth in this paper, we refer to these sets of data as the "shallow" and "deep" images, respectively. The boundaries of these images are indicated on images of σ Ori from the Digitized Sky Survey (DSS) in Figure 1.

The shallow IRAC images were collected on 2004 October 9 as a part of G. Fazio's IRAC Guaranteed Time Observations in *Spitzer* program 37 (Hernández et al. 2007a). Each IRAC mosaic for a given filter covered an area of 0.56 deg^2 ($0.7^\circ \times 0.8^\circ$). The overlapping area between the four filters was 0.5 deg^2 ($0.7^\circ \times 0.7^\circ$). At each cell in the mosaic and in each filter, IRAC obtained three 1 s exposures and three 26.8 s exposures.

The deep IRAC imaging was executed on 2006 September 28, 2007 March 30-31, and 2007 April 3-4 through *Spitzer* program 30395 (Scholz & Jayawardhana 2008). Instead of mosaics, 13 individual pointings were selected to encompass 18 candidate substellar members of σ Ori. Some of the images from these separate observations overlapped with each other, as shown in Figure 1. For each filter, a total area of 0.20 deg^2 was imaged. An area of 0.12 deg^2 was covered by all four bands. At each of the 13 pointings and in each of the 3.6, 4.5, and 5.8 μm filters, 12 images were obtained with exposure times of 96.8 s. For 8.0 μm, the number of images was doubled and the exposure times were 46.8 s.

The shallow images were processed with the Spitzer Science Center (SSC) S14.0.0 pipeline and the deep data were processed with the S14.4.0 and S15.3.0 pipelines. We combined the images produced by the SSC pipelines into mosaics and measured photometry for all point sources appearing in them using the methods described by Luhman et al. (2008). We selected a plate scale of $0.86''$ pixel^{-1} for the reduced IRAC mosaics, which is the native scale divided by $\sqrt{2}$. We used an aperture radius of 4 pixels when measuring photometry for most sources. Smaller apertures with radii of 2 or 3 pixels were applied to sources that were near other stars, which included the candidate brown dwarfs S Ori 25, S Ori 45, S Ori 47, S Ori 54, S Ori 65, S Ori 68, S Ori J053932.4−025220, S Ori J053949.5−023130, and S Ori J053929.4−024636. We also used a small aperture of 2 pixels for the new candidate member IRAC J05384729−0235194 (see § 3.2). The inner radius of the sky annulus was the same as the aperture radius in all cases. For the 4 pixel apertures at 5.8 and 8.0 μm, we selected relatively large widths of 6 pixels for the sky annuli to better measure the bright background emission at those wavelengths. In all other cases, the width of the annulus was 1 pixel. As noted above, some of the deep images overlapped with each other. For stars that were observed more than once during the spring observations, which spanned only a few days, we have adopted the average of the multiple measurements. Data obtained in both the fall and spring for a given star are presented separately. The completeness limits of the shallow images are 17.25, 17, 14.75, and 14 at 3.6, 4.5, 5.8, and 8.0 μm, respectively. The limits for the deep images are fainter by 0.75–1.25 mag. The quoted photometric errors include the Poisson errors in the source and background emission and the 2% uncertainty in the calibration of IRAC (Reach et al. 2005).

3. SEARCHING FOR NEW MEMBERS WITH DISKS

3.1. *Compilation of Probable Members*

In the first stage of our analysis of the disk population in σ Ori, we use our catalog of all sources detected by IRAC to search for new disk-bearing members of the cluster. We can use the IRAC data for known probable members of σ Ori to illustrate the typical colors of young stars with disks. To create a list of probable members, we begin with the sources that were considered by the other recent studies of disks in σ Ori (Hernández et al. 2007a; Caballero et al. 2007; Zapatero Osorio et al. 2007; Scholz & Jayawardhana 2008). We exclude sources from Hernández et al. (2007a) that are resolved as galaxies in images from the United Kingdom Infrared Telescope (UKIRT) Infrared Deep Sky Survey (UKIDSS, Lawrence et al. 2007)[6] or that have been spectroscopically classified as nonmembers by Kenyon et al. (2005) and Sacco et al. (2008). We also exclude stars that appear in the compilation of nonmembers from Caballero

---

[6] The UKIDSS project is described by Lawrence et al. (2007). UKIDSS uses the UKIRT Wide Field Camera (WFCAM, Casali et al. 2007) and a photometric system described by Hewett et al. (2006). The pipeline processing and science archive are described by Irwin et al. (in preparation) and Hambly et al. (2008). We have used data from the first data release, which is described in detail by Warren et al. (2007).

(2008) as well as source 668 from Hernández et al. (2007a), which is a galaxy according to spectroscopy from Caballero et al. (2008). S Ori 47 has been classified as a field dwarf rather than a cluster member based on gravity-sensitive absorption lines (McGovern et al. 2004), but we retain it in our compilation for the purposes of comparing our IRAC measurements to those from Caballero et al. (2007) and Scholz & Jayawardhana (2008). We add stars that have been spectroscopically confirmed as members by Scholz & Eislöffel (2004), Kenyon et al. (2005), Caballero (2006), and Sacco et al. (2008), the young stars associated with HH 446, Haro 5-22, and Haro 5-35, and σ Ori C and D. We also treat as members S Ori J053946.5−022423 and S Ori J053912.8−022453, which were identified as candidate members by Béjar et al. (2001) and exhibit evidence of disks in their IRAC colors. We present our IRAC measurements for this compilation of probable members in Table 1, which includes the source identifications from the 2MASS Point Source Catalog, Hernández et al. (2007a), Béjar et al. (1999, 2001), and the Henry Draper (HD) Catalog. We note that the evidence of membership varies significantly among the sources in Table 1, and that some of them are better characterized as possible members rather than probable members (e.g., S Ori 70, Burgasser et al. 2004).

3.2. *Candidate Members with IRAC Excesses*

A color-color diagram constructed from IRAC data is a convenient tool for efficiently identifying stars that exhibit IR excess emission indicative of disks. In Figure 2, we plot diagrams of $[3.6]-[4.5]$ versus $[5.8]-[8.0]$ for the probable members of σ Ori that are not saturated and that have photometric uncertainties less than 0.1 mag in all of the IRAC bands. The data from the shallow and deep images are shown separately. The stars in these diagrams reside near the origin or have significantly redder colors. These distinct populations are typical of young stellar populations (Hartmann et al. 2005) and represent stellar photospheres and stars with disks, respectively. In previous IRAC surveys of Taurus, Lupus, and Chamaeleon (Luhman et al. 2006, 2008; Allen et al. 2007), we have used color criteria of $[3.6]-[4.5] > 0.15$ and $[5.8]-[8.0] > 0.3$ to identify stars that might be new disk-bearing members. As shown in Figure 2, these colors also encompass most of the red population of probable members of σ Ori. Therefore, we have applied these criteria to the data in the bottom panels of Figure 2 to search for new members with disks. To further refine the sample of candidates, we have adopted an additional criterion of $[4.5]-[5.8] > 0.15$, which is satisfied by most stars with disks in σ Ori and in other clusters. The resulting candidates are plotted in the IRAC color-magnitude diagrams in Figure 3. Most of these sources are redder and fainter than the probable members of σ Ori, and thus are likely to be galaxies.

Seven of the candidates are in the list of uncertain members from Hernández et al. (2007a), consisting of sources 336, 361, 395, 614, 633, 916, and 950 from that study. Based on their faint magnitudes and red colors, sources 336, 361, 916, and 950 are either low-mass protostars or galaxies. Indeed, 950 is extended in K-band images from UKIDSS, indicating that it is probably a galaxy. Among the remaining candidates that are not in Hernández et al. (2007a), we have searched for sources that have colors similar to those of the probable members ($[3.6]-[8.0] < 2$), that are point sources in IRAC and UKIDSS images, and that are detected in the Z and K bands by UKIDSS. The five most promising candidate members are presented in Table 2. In addition to our IRAC data, we include photometry for these candidates measured from the 24 μm images from Hernández et al. (2007a). Two of the candidates, 2MASS J05383446−0253514 and 2MASS J05375398−0249545, are below the cluster sequence in the diagram of V versus $V-J$ from Hernández et al. (2007a), which may indicate that they are seen in scattered light (e.g., edge-on disks) if they are bona fide members. The latter candidate also appears near the lower edge of the cluster sequence in V versus $V-I$ (Sherry et al. 2004). Although it is underluminous in V versus $V-J$, 2MASS J05383446−0253514 does appear within the cluster sequence in I versus $I-K$ (Caballero 2008), probably because of K-band excess emission. 2MASS J05375398−0249545 was detected in an Hα objective prism survey (Weaver & Babcock 2004), providing additional evidence that it could be a young star. Based on their IRAC magnitudes, these two candidates should have masses of $\sim 0.5\ M_\odot$ if they are cluster members. The candidates IRAC J05391066−0229238 and IRAC J05384459−0225433 have much fainter magnitudes that are indicative of brown dwarfs. The former underwent significant variability in all IRAC bands between the shallow and deep images. The final candidate, IRAC J05384729−0235194, exhibits very red IRAC colors that are consistent with either a protostar or a galaxy. This object is only $6''$ from source 726 from Hernández et al. (2007a) and is $1'$ from σ Ori. Its close proximity to the center of the cluster and its relatively bright magnitude compared to most galaxies (see Figure 3) tend to favor membership.

4. DISK POPULATION

4.1. *Identifying the Members that have Disks*

To characterize the disk population among low-mass members of the σ Ori cluster, we begin by using the IRAC data to identify the probable members that are likely to have disks. Because a disk produces greater excess emission above a stellar photosphere at longer wavelengths, we select the 5.8 and 8.0 μm bands of IRAC for measuring disk emission. To construct colors with these data that can be used for measuring excess emission, we combine them with the IRAC data at 3.6 μm, which are available for nearly all of the probable members that are measured at 5.8 and 8.0 μm. Measuring excess emission from $[3.6]-[5.8]$ and $[3.6]-[8.0]$ requires estimates of the intrinsic colors for stellar photospheres and their dependence on spectral type. We can obtain these estimates by plotting the colors as a function of magnitude, which is a reasonable proxy for spectral type in a sample of stars that are roughly coeval and have low reddening ($A_V < 5$). In Figure 4, we show color-magnitude diagrams of this kind for the probable members of σ Ori. The stars that lack disk emission in the IRAC bands form a well-defined sequence near colors of zero. In both $[3.6]-[5.8]$ and $[3.6]-[8.0]$, the sequence becomes slightly redder with fainter magnitudes, which reflects the dependence on spectral type. For comparison, we include in

Figure 4 a fit to the sequence of diskless members of the Chamaeleon I star forming region (Luhman et al. 2008). The averages of [3.6]−[5.8] and [3.6]−[8.0] for all stars detected by IRAC toward Chamaeleon I, which are mostly background stars, are 0.05 mag greater than the average values in σ Ori, which is consistent with the difference in average extinction for the two regions. Therefore, we have dereddened the sequence for Chamaeleon I accordingly in each color in Figure 4. We have also corrected for differences in ages and distances by adding 1.6 mag to the Chamaeleon I fits, which is the average difference between the two cluster sequences in [3.6] versus spectral type. After applying these color and magnitude offsets, the sequence for Chamaeleon I agrees well that of σ Ori.

To identify members that exhibit significant excess emission in [3.6] − [5.8] and [3.6] − [8.0] and thus are likely to have disks, we must properly account for errors and biases in the photometric measurements, particularly near the detection limits of the data. As mentioned in § 2, our reported errors include Poisson errors and a 2% uncertainty in the calibration of IRAC. However, additional systematic errors are also present, such as location-dependent variations in the calibration. The precision with which we can measure excess emission is also limited by the intrinsic scatter of photospheric colors among cluster members at a given magnitude. Fortunately, in a well-populated sample of stars, the sum of these uncertainties is reflected in the spread in colors at a given magnitude among diskless cluster members. Therefore, we can use the widths of the sequences in Figure 4 to define color thresholds for identifying sources that have significant color excesses. For the shallow data, we have computed the standard deviation of colors in each sequence as a function of magnitude for the range of magnitudes in which the sequences are sufficiently populated ([3.6] < 14.25). Stars that are redder than the 2 σ thresholds are classified as having disks. Because the sequences in the deep images contain fewer stars, we adopt for those data the thresholds from the shallow images after shifting them to fainter magnitudes by 0.75 mag, which is the difference in completeness limits between the shallow and deep [8.0] data. The classifications produced by [3.6] − [5.8] and [3.6] − [8.0] agree for 99.5% of the sources. For the remaining 0.5% of the sample, we adopt the classifications from [3.6] − [8.0] since this color is more sensitive to disk emission. Our classifications apply only to inner disks that are capable of producing emission at 8 μm. A few of the stars that exhibit photospheric colors in the IRAC bands may have disks with inner holes that produce excess emission only at longer wavelengths.

At the faintest magnitudes in Figure 4, the photospheric sequences are too sparsely populated for reliable measurements of their widths. To classify sources that are fainter than the 2 σ thresholds, we rely on a comparison of the Chamaeleon I sequence to the Poisson errors, which are indicated in Figure 4 for the brown dwarf candidates that were considered by Zapatero Osorio et al. (2007) and Scholz & Jayawardhana (2008). For the intrinsic photospheric colors of the faintest source, S Ori 70, we adopt the average colors measured for field dwarfs (Patten et al. 2006) that have a similar spectral type (T6; Burgasser et al. 2006). At the faintest magnitudes, the Poisson errors should be much larger than the systematic errors from the location-dependent variations in the calibration and the intrinsic scatter in photospheric colors. However, there remains an additional source of error that must be considered. In background-limited images, a faint source is preferentially detected if upward fluctuations of the background noise bring it above the detection limit. As a result, the fluxes measured for sources with very low signal-to-noise ratios (SNRs) are systematically overestimated. For instance, fluxes near the detection limits of the Two-Micron All-Sky Survey (2MASS; Skrutskie et al. 2006) are overestimated by an average of $\sim$ 0.4 mag (Beichman et al. 2003). At SNR< 5, this bias and the Poisson errors are comparable to color excesses from disks, making such measurements difficult to use in our analysis. By comparing the color excesses relative to the Chamaeleon I sequence to the sizes of the Poisson errors and ignoring the data at SNR< 5, we have identified the faint sources in Figure 4 that are likely to have disks. These classifications for the brown dwarf candidates considered by Zapatero Osorio et al. (2007) and Scholz & Jayawardhana (2008) are presented in Table 3. The available data at 5.8 and 8.0 μm offer insufficient SNRs for classifying six of the sources in Table 3 as well as S Ori J053844.5−025512, S Ori J053932.4−025220, S Ori J053929.4−024636, S Ori J053944.5−025959, and S Ori J054007.0−023604.

Based on our classifications in Table 3, S Ori 60 is the faintest object in σ Ori that exhibits significant evidence of a disk. If we adopt a distance modulus of 7.65 (§ 4.2) and photometry from UKIDSS[7], then S Ori 60 has $M_J = 11.3$, $M_H = 10.5$, and $M_K = 9.9$. In comparison, the least luminous disk-bearing brown dwarf found in other young clusters is Cha J11070768−7626326, which has slightly fainter magnitudes of $M_J = 11.56$, $M_H = 10.75$, and $M_K = 9.86$ (Luhman et al. 2008). The faintest σ Ori source that has marginal excess emission is S Ori 66. It is $\sim$ 0.8 mag fainter than S Ori 60 according to UKIDSS.

4.2. *Disk Fraction*

We now use the IRAC classifications derived in the previous section to compute the disk fraction among the probable members of σ Ori. We wish to measure the dependence of this disk fraction on stellar mass. Ideally, the masses would be estimated by combining spectral types and luminosities with theoretical evolutionary models. However, spectral classifications are unavailable for many of the probable members of σ Ori. Therefore, we adopt M_J as a proxy for stellar mass, as done by Hernández et al. (2007a). We adopt the J measurements from 2MASS and UKIDSS when possible. For the few sources that do not have photometric errors less than 0.2 mag in these surveys, we use the J data from Béjar et al. (2001), Martín et al. (2001), Caballero et al. (2007), and Zapatero Osorio et al. (2008). We assume that the probable members of σ Ori have negligible extinctions at J. For comparison to σ Ori, we also consider other clusters in which disk fractions that have been measured for both stars and brown dwarfs using IRAC data. Therefore, we have recomputed the disk fractions for IC 348 and Chamaeleon I from Muench et al.

---

[7] We adopt the UKIDSS magnitudes measured with an aperture radius of 1″ (Dye et al. 2006).

(2007) and Luhman et al. (2008) as a function of M_J using the J-band data from 2MASS, Luhman et al. (2003), and Luhman (2007) and extinction estimates from Muench et al. (2007) and Luhman (2007). We adopt distance moduli of 7.5 and 6.05 for IC 348 and Chamaeleon I, respectively (Herbig 1998; Luhman 2008). To enable a meaningful comparison of the disk fractions of these clusters, M_J must be corrected for differences in age so that a given value of M_J corresponds to the same mass in each cluster. For the adopted distances, IC 348 and Chamaeleon I do have similar ages (Luhman 2007). In order to compare the disk fraction in σ Ori to those of the other two clusters in a manner that is as age-independent as possible, we adopt a distance modulus that is the sum of the distance modulus for Chamaeleon I and the difference of the apparent magnitudes as a function of spectral type between Chamaeleon I and σ Ori (§ 4.1), corresponding to a value of 7.65 (340 pc). In other words, we are adopting a distance for σ Ori that gives it the same age as IC 348 and Chamaeleon I. Similar distances have been adopted in some previous studies of σ Ori (Béjar et al. 2001; Oliveira et al. 2002) while values near 440 pc have been used elsewhere (Sherry et al. 2004; Hernández et al. 2007a). The larger distances would imply that σ Ori is younger than IC 348 and Chamaeleon I.

Six sources in σ Ori are beyond the faint magnitude limit that we have selected for plotting the disk fractions ($M_J > 12$, $J > 19.65$), four of which are too faint for a useful constraint on excess emission at 5.8 or 8.0 μm. In the faintest bin of the disk fraction for σ Ori ($M_J = 10$–12), two objects have tentative detections of excesses and two others are too faint for classifications. For the purposes of this work, we assume that the disks are present in the former but not the latter.

The disk fractions versus M_J for σ Ori, IC 348, and Chamaeleon I are listed in Table 4 and are plotted in Figure 5. For $M_J > 4$ ($M \lesssim 0.5\ M_\odot$), the three regions exhibit similar disk fractions. In each case, $\sim 40\%$ of low-mass stars (0.08–0.5 $M_\odot$) and ~ 50–60% of brown dwarfs (0.01–0.08 $M_\odot$) have disks. Meanwhile, the disk fractions at $M_J < 4$ range from $\sim 30\%$ in σ Ori to $\sim 65\%$ in Chamaeleon I. Since these clusters have similar ages, these data suggest that the lifetimes of disks around solar-mass stars increase from σ Ori to IC 348 to Chamaeleon I.

We note a few caveats regarding the disk fractions in Figure 5. A given M_J magnitude corresponds to a larger range of stellar masses than a given spectral type. In addition, the near-IR magnitudes of stars with edge-on disks are typically much fainter than most other cluster members at the same spectral type and color. As a result, a star with an edge-on disk can appear at a magnitude bin corresponding to substellar masses in a plot of disk fraction versus M_J. Meanwhile, stars with disks can have systematically brighter J magnitudes than diskless stars because of disk emission in this band, which would cause these sources to appear in magnitude bins that are too bright. Because of these effects, the disk fractions versus M_J in Figure 5 are smoothed versions of the disk fractions versus mass or spectral type. For instance, the difference between the disk fractions of Chamaeleon I and IC 348 at high masses is more pronounced when they are plotted as a function of spectral type or mass (Luhman et al. 2008). Finally, unlike the samples for IC 348 and Chamaeleon I, some of the sources included in the disk fraction for σ Ori lack definitive evidence of membership. Thus, our estimate of the disk fraction for σ Ori is may be underestimated. Indeed, Sacco et al. (2008) obtained spectra of 23% of the probable members identified by Hernández et al. (2007a) and found a field star contamination rate of 20%. Because of the spectral classifications that are now available from Sacco et al. (2008), the contamination of our current sample of probable members should be lower.

4.3. *Comparison to Previous Studies*

As discussed in § 1, five previous studies have used IRAC data to investigate the disk population among brown dwarfs in the σ Ori cluster (Hernández et al. 2007a; Caballero et al. 2007; Zapatero Osorio et al. 2007, 2008; Scholz & Jayawardhana 2008). To compare the results from those studies and this work, we begin by plotting the differences between the previously reported IRAC magnitudes and our measurements for the shallow and deep images in Figures 6 and 7, respectively. The differences in photometry are more than 0.1 mag in many cases and reach as high as 1–2 mag. Most of the measurements from Caballero et al. (2007) and Zapatero Osorio et al. (2007) agree with our data within their quoted errors, but not within our errors. Roughly half of the measurements in common between this work and Scholz & Jayawardhana (2008) differ by an amount that is larger than the errors from either study. Our formal errors are much lower than the ones reported by Caballero et al. (2007), Zapatero Osorio et al. (2007), and Scholz & Jayawardhana (2008), but this in itself does not demonstrate that our measurements are truly more accurate. To test the relative accuracies of these sets of photometry, we compare them in color-color diagrams in Figure 8. For both the shallow and deep images, our colors are more tightly clustered into two groups (diskless and disk-bearing stars) than the colors in the other studies, which suggests that our colors indeed have lower errors. A similar result was found by Luhman et al. (2008) in a comparison of their IRAC data to measurements from Damjanov et al. (2007) in Chamaeleon I.

We now compare our classifications of the IRAC data in σ Ori to those derived in previous work. The classifications from Caballero et al. (2007) and our earlier survey of σ Ori (Hernández et al. 2007a) agreed for the 25 sources that were considered by both studies, which were predominantly low-mass stars. The samples from Zapatero Osorio et al. (2007) and Scholz & Jayawardhana (2008) consisted of fainter brown dwarf candidates. These sources are listed in Table 3, where we compare the classifications from Caballero et al. (2007), Zapatero Osorio et al. (2007), Scholz & Jayawardhana (2008), and this work. The four studies differ significantly in the sources that are found to exhibit excess emission from disks. Zapatero Osorio et al. (2007) reported excess emission at 5.8 or 8.0 μm in the shallow images for seven candidate brown dwarfs. However, we find that six of these sources have SNRs in those data that are too low for useful photometry, as illustrated in Figure 9. Four of these six candidates are within the deep images from Scholz & Jayawardhana (2008), which provide much better detections. Using the deep data, we measure significant excess emission for S Ori 56 and S Ori 60 and marginal excesses for S Ori 55 and S Ori 58. Scholz & Jayawardhana (2008) derived the same classifications for S Ori 58 and S Ori 60, but concluded that the other two sources lack disks. The seventh object from Zapatero Osorio et al. (2007), S Ori 54, is too close ($5''$) to a red galaxy for an accurate measurement at 8 μm using either the shallow or deep images. We do not find excess emission for this source at 5.8 μm, where the contrast relative to the galaxy is more favorable.

The three remaining sources Table 3 that have discrepant classifications are S Ori 65, S Ori 70, and S Ori J053949.5−023130. Scholz & Jayawardhana (2008) found significant excess emission for S Ori 65 and S Ori J053949.5−023130 at 5.8 and 8.0 μm from their deep images. However, our measurements at 5.8 μm do not exhibit excesses and the detections at 8.0 μm are too weak to be useful (see Figure 10). Although S Ori 65 has $[3.6]-[8.0]\sim 1$ in our data, which implies an excess, the photometry at 8.0 μm has an error of 0.4 mag, corresponding to SNR~ 3. As discussed in § 4.1, measurements with such low values of SNR are subject to flux overestimation. To illustrate that color excesses with such large errors are not significant, we note that another source with comparable errors at 8.0 μm, S Ori 50, has a nominal color that is unphysical for stellar sources ($[3.6]-[8.0]\sim -0.9$). Finally, based on their deep images, Scholz & Jayawardhana (2008) suggested that S Ori 70 may have excesses in $[3.6]-[4.5]$ and $[3.6]-[5.8]$ relative to typical colors of T6 dwarfs, which they attributed to either a disk or low surface gravity and cited as possible evidence of youth and membership in the σ Ori cluster. Zapatero Osorio et al. (2008) arrived at a similar conclusion through a measurement of $[3.6]-[4.5]$ from the shallow images. We measure $[3.6]-[4.5]=1.56\pm0.07$ and $[3.6]-[5.8]=1.75\pm0.21$ from the deep data. In comparison, the colors of field dwarfs between T6 and T6.5 range from 0.93 to 1.46 in $[3.6]-[4.5]$ and from 0.87 to 1.19 in $[3.6]-[5.8]$ (Patten et al. 2006). Thus, we find that S Ori 70 is only slightly redder in $[3.6]-[4.5]$ than the reddest field dwarfs. Given the low SNR at 5.8 μm (see Figure 10), the apparent excess in this band could be caused by a combination of Poisson errors and flux overestimation.

5. CONCLUSIONS

We have investigated the disk population among stars and brown dwarfs in the σ Ori cluster using mid-IR images obtained with IRAC onboard the *Spitzer Space Telescope*. For this study, we have employed all available IRAC data for σ Ori, which consist of shallow images (80.4 s) from Hernández et al. (2007a) and deep images ($\sim$ 1100 s) from Scholz & Jayawardhana (2008). We measured photometry for all sources detected in these images and searched the resulting data for new members of the cluster based on the red colors that are expected from stars with disks. The five most promising candidates have colors and magnitudes that are suggestive of edge-on disks, brown dwarfs, and a low-mass protostar. We then examined the IRAC colors for $\sim$ 300 probable cluster members found in previous studies and identified the ones that are likely to have disks. In doing so, we have attempted to fully account for the errors and biases in the photometry of the brown dwarf candidates (e.g., flux overestimation at low SNR; Beichman et al. 2003), some of which are near the detection limits of the IRAC data. S Ori 60 is the faintest candidate member of σ Ori that exhibits significant IR excess emission. This object is comparable in luminosity to the faintest brown dwarf that shows evidence of a disk in other star-forming regions (Luhman et al. 2008). Using our classifications of the IRAC data, we computed the disk fraction as a function of M_J, which acts as a proxy for stellar mass. The disk fractions for low-mass stars (0.08–0.5 $M_\odot$) and brown dwarfs (0.01–0.08 $M_\odot$) are $\sim$ 40% and $\sim$ 60%, respectively, which are similar to the disk fractions derived from IRAC surveys of two other clusters near the same age, IC 348 and Chamaeleon I (Muench et al. 2007; Luhman et al. 2008).

Although our disk fraction for brown dwarfs in σ Ori is similar to other estimates based on the IRAC data considered here (Caballero et al. 2007; Zapatero Osorio et al. 2007; Scholz & Jayawardhana 2008), our photometric measurements and our classifications of the IRAC colors (disk vs. no disk) differ significantly from those in previous studies. For instance, Zapatero Osorio et al. (2008) and Scholz & Jayawardhana (2008) suggested that the T dwarf S Ori 70 may exhibit color excesses in $[3.6]-[4.5]$ and $[3.6]-[5.8]$ relative to field dwarfs at the same spectral type. They cited these non-standard colors as possible evidence of youth, which would indicate that S Ori 70 is a cluster member rather than a field dwarf. However, we have found that this object is not significantly redder in $[3.6]-[4.5]$ than field dwarfs and that its SNR may be too low at 5.8 μm for a useful measurement of $[3.6]-[5.8]$. Therefore, we conclude that the IRAC data do not provide firm constraints on the membership of S Ori 70.

K. L. was supported by grant AST-0544588 from the National Science Foundation. This publication has made use of NASA's Astrophysics Data System Bibliographic Services, the SIMBAD database, operated at CDS in Strasbourg, France, and data products from 2MASS and DSS. 2MASS is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. DSS was produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166.

REFERENCES

Allen, L. E., et al. 2004, ApJS, 154, 363
Allen, P. R., et al. 2007, ApJ, 655, 1095
Barrado y Navascués, D., Zapatero Osorio, M. R., Martín, E. L., Béjar, V. J. S., Rebolo, R., & Mundt, R. 2002, A&A, 393, L85
Barrado y Navascués, D., et al. 2007, ApJ, 664, 481
Beichman, C. A., Cutri, R., Jarrett, T., Stiening, R., & Skrutskie, M. 2003, AJ, 125, 2521
Béjar, V. J. S., Zapatero Osorio, M. R., & Rebolo, R. 1999, ApJ, 521, 671
Béjar, V. J. S., et al. 2001, ApJ, 556, 830

Burgasser, A. J., Kirkpatrick, J. D., Reid, I. N., Brown, M. E., Miskey, C. L., & Gizis, J. E. 2003, ApJ, 586, 512
Burgasser, A. J., Kirkpatrick, J. D., McGovern, M. R., McLean, I. S., Prato, L., & Reid, I. N. 2004, ApJ, 604, 827
Burgasser, A. J., et al. 2006, ApJ, 637, 1067
Caballero, J. A. 2006, PhD thesis, Univ. de La Laguna, Spain
Caballero, J. A. 2008, A&A, 478, 667
Caballero, J. A., et al. 2007, A&A, 470, 903
Caballero, J. A., et al. 2008, A&A, in press
Carpenter, J. M., J. M., Mamajek, E. E., Hillenbrand, L. A., Meyer, M. R. ApJ, 651, L49
Casali, M., et al. 2007, A&A, 467, 777
Dahm, S. E., & Hillenbrand, L. A. 2007, AJ, 133, 2072
Damjanov, I., Jayawardhana, R., Scholz, A., Ahmic, M., Nguyen, D. C., Brandeker, A., & van Kerkwijk, M. H. 2007, ApJ, 670, 1337
Dye, S., et al. 2006, MNRAS, 372, 1227
Fazio, G. G., et al. 2004, ApJS, 154, 10
Flaherty, K. M., Pipher, J. L., Megeath, S. T., Winston, E. M., Gutermuth, R. A., Muzerolle, J., Allen, L. E., & Fazio, G. G. 2007, ApJ, 663, 1069
Guieu, S., et al. 2007, A&A, 465, 855
Gutermuth, R. A., Megeath, S. T., Muzerolle, J., Allen, L. E., Pipher, J. L., Myers, P. C., & Fazio, G. G. 2004, ApJS, 154, 374
Haisch, K. E., Lada, E. A., & Lada, C. J. 2001, ApJ, 553, L153
Hambly, N. C., et al. 2008, MNRAS, 384, 637
Hartmann, L., Megeath, S. T., Allen, L., Luhman, K., Calvet, N., D'Alessio, P., Franco-Hernandez, R., & Fazio, G. 2005, ApJ, 629, 881
Herbig, G. H. 1998, ApJ, 497, 736
Hernández, J., Calvet, N., Hartmann, L., Briceño, C., Sicilia-Aguilar, A., & Berlind, P. 2005, AJ, 129, 856
Hernández, J., Hartmann, L., Calvet, N., Jeffries, R. D., Gutermuth, R., Muzerolle, J., & Stauffer, J. 2008, ApJ, in press
Hernández, J., et al. 2007a, ApJ, 662, 1067
Hernández, J., et al. 2007b, ApJ, 671, 1784
Hewett, P. C., Warren S. J., Leggett S. K., & Hodgkin S. L., 2006, MNRAS, 367, 545
Kenyon, M. J., Jeffries, R. D., Naylor, T., Oliveira, J. M., & Maxted, P. F. L. 2005, MNRAS, 356, 89
Kenyon, S. J., & Hartmann, L. 1995, ApJS, 101, 117
Lada, C. J., et al. 2006, AJ, 131, 1574
Lawrence, A., et al. 2007, MNRAS, 379, 1599
Luhman, K. L. 2007, ApJS, 173, 104
Luhman, K. L. 2008, Handbook of Star Forming Regions, ASP Conference Series, in press
Luhman, K. L., Stauffer, J. R., Muench, A. A., Rieke, G. H., Lada, E. A., Bouvier, J., & Lada, C. J. 2003, ApJ, 593, 1093
Luhman, K. L., Whitney, B. A., Meade, M. R., Babler, B. L., Indebetouw, R., Bracker, S., & Churchwell, E. B. 2006, ApJ, 647, 1180
Luhman, K. L., et al. 2005, ApJ, 631, L69
Luhman, K. L., et al. 2008, ApJ, 675, 1375
Martín, E. L., & Zapatero Osorio, M. R. 2003, ApJ, 593, L113
Martín, E. L., Zapatero Osorio, M. R., Barrado y Navascués, D., Béjar, V. J. S., & Rebolo, R. 2001, ApJ, 558, L117
McGovern, M. R., Kirkpatrick, J. D., McLean, I. S., Burgasser, A. J., Prato, L., & Lowrance, P. J. 2004, ApJ, 600, 1020
Megeath, S. T., Hartmann, L., Luhman, K. L., & Fazio, G. G. 2005, ApJ, 646, 1215
Megeath, S. T., et al. 2004, ApJS, 154, 367
Muench, A. A., Lada, C. J., Luhman, K. L., Muzerolle, J., & Young, E. 2007, AJ, 134, 411
Muzerolle, J., et al. 2004, ApJS, 154, 379
Oliveira, J. M., Jeffries, R. D., Kenyon, M. J., Thompson, S. A., & Naylor, T. 2002, A&A, 382, L22
Patten, B. M., et al. 2006, ApJ, 651, 502
Reach, W. T., et al. 2005, PASP, 117, 978
Sacco, G. G., Franciosini, E., Randich, S., & Pallavicini, R. 2008, A&A, in press
Scholz, A., & Eislöffel, J. 2004, A&A, 419, 249
Scholz, A., & Jayawardhana, R. 2008, ApJ, 672, L49
Sherry, W. H., Walter, F. M., & Wolk, S. J. 2004, AJ, 128, 2316
Sicilia-Aguilar, A., et al. 2006, ApJ, 638, 897
Skrutskie, M., et al. 2006, AJ, 131, 1163
Warren, S. J., et al. 2007, MNRAS, 375, 213
Weaver, W. B., & Babcock, A. 2004, PASP, 116, 1035
Werner, M. W., et al. 2004, ApJS, 154, 1
Zapatero Osorio, M. R., Béjar, V. J. S., Martín, E. L., Rebolo, R., Barrado y Navascués, D., Mundt, R., Eislöffel, J., & Caballero, J. A. 2002a, ApJ, 578, 536
Zapatero Osorio, M. R., Béjar, V. J. S., Pavlenko, Ya., Rebolo, R., Allende Prieto, C., Martín, E. L., & Garca López, R. J. 2002b, A&A, 384, 937
Zapatero Osorio, M. R., et al. 2007, A&A, 472, L9
Zapatero Osorio, M. R., et al. 2008, A&A, 477, 895

TABLE 1
Spitzer Photometry for Probable Members of σ Ori

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
2MASS J05371537−0230534	···	sat	sat	10.69±0.03	10.79±0.03	2006 Sep 28
2MASS J05371869−0240218	SO 9	11.01±0.02	10.98±0.02	10.81±0.03	10.94±0.03	2004 Oct 9
		out	11.01±0.02	out	10.94±0.03	2006 Sep 28
2MASS J05372306−0232465	SO 27	13.02±0.02	12.95±0.02	12.90±0.04	13.03±0.04	2004 Oct 9
		13.03±0.02	12.95±0.02	12.89±0.03	12.90±0.03	2006 Sep 28
IRAC J05372470−0231523	S Ori 66	17.43±0.06	16.82±0.08	···	···	2004 Oct 9
		17.21±0.03	16.80±0.04	16.27±0.09	16.26±0.21	2006 Sep 28
IRAC J05372587−0234320	S Ori 55	16.49±0.04	16.23±0.05	···	···	2004 Oct 9
		16.44±0.02	16.14±0.02	15.93±0.07	15.51±0.11	2006 Sep 28
2MASS J05372761−0257100	···	out	13.88±0.02	out	13.81±0.06	2004 Oct 9
2MASS J05372806−0236065	SO 59	12.66±0.02	12.58±0.02	12.51±0.03	12.60±0.04	2004 Oct 9
		12.72±0.02	12.53±0.02	12.63±0.03	12.59±0.03	2006 Sep 28
2MASS J05372831−0224182	SO 60	12.88±0.02	12.81±0.02	12.79±0.03	12.83±0.04	2004 Oct 9
		12.90±0.02	out	12.76±0.03	out	2006 Sep 28
2MASS J05373094−0223427	SO 73	10.90±0.02	10.43±0.02	10.04±0.03	9.12±0.03	2004 Oct 9
		sat	out	10.15±0.03	out	2006 Sep 28
2MASS J05373153−0224269	SO 77	10.99±0.02	10.96±0.02	10.93±0.03	10.95±0.03	2004 Oct 9
		sat	out	10.86±0.03	out	2006 Sep 28
2MASS J05373187−0245184	SO 82,HD 294275	9.22±0.02	9.20±0.02	9.16±0.03	9.22±0.03	2004 Oct 9
2MASS J05373645−0214375	SO 115	10.73±0.02	10.71±0.02	10.67±0.03	10.72±0.03	2004 Oct 9
2MASS J05373648−0241567	SO 116,S Ori 40	14.19±0.02	14.03±0.02	14.00±0.04	14.03±0.07	2004 Oct 9
		out	14.00±0.02	out	13.97±0.04	2006 Sep 28
2MASS J05373666−0234003	SO 117	11.83±0.02	11.82±0.02	11.81±0.03	11.79±0.03	2004 Oct 9
		sat	11.74±0.00	11.80±0.03	11.74±0.03	2006 Sep 28
2MASS J05373784−0245442	SO 123	11.49±0.02	11.47±0.02	11.42±0.03	11.46±0.03	2004 Oct 9
2MASS J05374248−0256358	···	out	14.51±0.02	out	14.26±0.08	2004 Oct 9
2MASS J05374491−0229573	SO 164	9.89±0.02	9.88±0.02	9.89±0.03	9.91±0.03	2004 Oct 9
2MASS J05374527−0228521	SO 165	12.67±0.02	12.55±0.02	12.56±0.03	12.53±0.04	2004 Oct 9
2MASS J05374557−0229585	···	10.47±0.02	10.45±0.02	10.47±0.03	10.47±0.03	2004 Oct 9
2MASS J05374963−0236182	SO 194	10.11±0.02	10.15±0.02	10.04±0.03	10.12±0.03	2004 Oct 9
2MASS J05375110−0226074	SO 209,S Ori 23	13.66±0.02	13.51±0.02	13.63±0.04	13.52±0.06	2004 Oct 9
2MASS J05375161−0235257	SO 214	10.78±0.02	10.79±0.02	10.80±0.03	10.73±0.03	2004 Oct 9
2MASS J05375206−0236046	SO 219	13.95±0.02	13.79±0.02	13.76±0.04	13.73±0.06	2004 Oct 9
2MASS J05375210−0256551	···	out	12.12±0.02	out	12.10±0.03	2004 Oct 9
2MASS J05375220−0233379	SO 220	11.82±0.02	11.83±0.02	11.81±0.03	11.77±0.03	2004 Oct 9
2MASS J05375303−0233344	SO 229	9.45±0.02	9.46±0.02	9.44±0.03	9.43±0.03	2004 Oct 9
2MASS J05375404−0244407	SO 240	11.90±0.02	11.87±0.02	11.85±0.03	11.83±0.03	2004 Oct 9
2MASS J05375440−0239298	SO 243	8.53±0.02	8.69±0.02	8.55±0.03	8.51±0.03	2004 Oct 9
2MASS J05375445−0243378	SO 244	8.98±0.02	9.05±0.02	8.99±0.03	8.98±0.03	2004 Oct 9
2MASS J05375452−0258265	···	out	11.95±0.02	out	11.96±0.03	2004 Oct 9
2MASS J05375486−0241092	SO 247	12.04±0.02	11.79±0.02	11.45±0.03	10.84±0.03	2004 Oct 9
2MASS J05375512−0227362	SO 251	12.45±0.02	12.34±0.02	12.33±0.03	12.48±0.04	2004 Oct 9
		12.40±0.02	out	12.30±0.03	out	2006 Sep 28
2MASS J05375559−0233053	SO 254,S Ori 35	13.66±0.02	13.32±0.02	12.98±0.04	12.34±0.04	2004 Oct 9
2MASS J05375574−0224337	SO 256,S Ori 24	13.79±0.02	13.71±0.02	13.63±0.04	13.73±0.07	2004 Oct 9
2MASS J05375647−0243574	SO 260	11.92±0.02	11.94±0.02	11.88±0.03	11.88±0.03	2004 Oct 9
2MASS J05375745−0238444	SO 271,S Ori 12	12.90±0.02	12.74±0.02	12.59±0.03	12.03±0.03	2004 Oct 9
2MASS J05375840−0241262	SO 283	12.06±0.02	11.95±0.02	11.91±0.03	11.99±0.03	2004 Oct 9
2MASS J05380055−0245097	SO 297	11.57±0.02	11.46±0.02	11.48±0.03	11.45±0.03	2004 Oct 9
2MASS J05380097−0226079	SO 299	11.74±0.02	11.73±0.02	11.68±0.03	11.37±0.03	2004 Oct 9
2MASS J05380107−0245379	SO 300	10.13±0.02	9.69±0.02	9.30±0.03	8.66±0.03	2004 Oct 9
2MASS J05380167−0225527	SO 302	11.88±0.02	11.83±0.02	11.80±0.03	11.83±0.03	2004 Oct 9
2MASS J05380421−0219057	SO 320	10.71±0.02	10.75±0.02	10.76±0.03	10.71±0.03	2004 Oct 9
		sat	10.81±0.00	10.78±0.03	10.71±0.03	2007 Apr 3
2MASS J05380552−0235571	SO 327,S Ori J053805.5-023557	13.74±0.02	13.43±0.02	13.13±0.04	12.45±0.04	2004 Oct 9
		13.72±0.02	13.42±0.02	13.07±0.03	12.47±0.03	2006 Sep 28
		out	13.46±0.02	out	12.51±0.03	2007 Mar 31
2MASS J05380649−0228494	SO 338	9.70±0.02	9.66±0.02	9.66±0.03	9.68±0.03	2004 Oct 9
		sat	out	9.66±0.03	out	2006 Sep 28
		sat	out	9.65±0.03	out	2007 Apr 3
2MASS J05380674−0230227	SO 341	9.77±0.02	9.46±0.02	9.19±0.03	8.45±0.03	2004 Oct 9
		sat	out	9.26±0.03	out	2006 Sep 28
		sat	out	9.02±0.03	out	2007 Apr 3
2MASS J05380784−0231314	SO 352	9.62±0.02	9.61±0.02	9.57±0.03	9.59±0.03	2004 Oct 9
		sat	out	9.65±0.03	out	2006 Sep 28
		sat	sat	out	9.58±0.03	2007 Mar 31/Apr 3
2MASS J05380826−0235562	SO 362	10.71±0.02	10.44±0.02	10.11±0.03	9.11±0.03	2004 Oct 9
		sat	sat	10.10±0.03	9.03±0.03	2006 Sep 28
		out	sat	out	8.95±0.03	2007 Mar 31
2MASS J05380897−0220109	SO 366	12.08±0.02	12.05±0.02	12.03±0.03	12.04±0.03	2004 Oct 9
		12.10±0.02	12.03±0.02	12.04±0.03	11.99±0.03	2007 Apr 3
2MASS J05380994−0251377	SO 374	10.70±0.02	10.37±0.02	10.13±0.03	9.50±0.03	2004 Oct 9
2MASS J05381012−0254506	SO 377	13.07±0.02	12.91±0.02	12.99±0.04	12.92±0.05	2004 Oct 9
IRAC J05381013−0236261	S Ori 70	18.38±0.14	17.04±0.09	···	···	2004 Oct 9
		18.76±0.06	17.20±0.03	17.01±0.20	16.62±0.51	2006 Sep 28

TABLE 1 — *Continued*

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
		out	17.44±0.10	out	···	2007 Mar 31
2MASS J05381064−0232574	SO 379	10.51±0.02	10.54±0.02	10.48±0.03	10.46±0.03	2004 Oct 9
		out	sat	out	10.48±0.03	2007 Mar 31
2MASS J05381175−0245012	SO 385	9.33±0.02	9.48±0.02	9.38±0.03	9.38±0.03	2004 Oct 9
		out	sat	out	9.32±0.03	2006 Sep 28
		sat	out	9.41±0.03	out	2007 Mar 31
2MASS J05381223−0218124	SO 389	11.76±0.02	11.82±0.02	11.77±0.03	11.78±0.03	2004 Oct 9
2MASS J05381239−0219387	SO 392	13.91±0.02	13.55±0.02	13.38±0.04	12.73±0.04	2004 Oct 9
		13.90±0.02	13.53±0.02	13.33±0.03	12.83±0.03	2007 Apr 3
IRAC J05381262−0221375	S Ori 67	18.38±0.15	···	···	···	2004 Oct 9
		18.12±0.04	18.10±0.09	···	···	2007 Apr 3
2MASS J05381279−0212266	···	12.92±0.02	out	12.29±0.03	out	2004 Oct 9
		out	12.55±0.02	out	11.56±0.03	2007 Apr 3
2MASS J05381315−0245509	SO 396	10.06±0.02	9.67±0.02	9.46±0.03	8.90±0.03	2004 Oct 9
		out	sat	out	8.87±0.03	2006 Sep 28
		sat	sat	9.44±0.03	8.79±0.03	2007 Mar 31/Apr 3
2MASS J05381319−0226088	SO 397	11.17±0.02	10.95±0.02	10.68±0.03	9.99±0.03	2004 Oct 9
		sat	out	10.75±0.03	out	2007 Apr 3
2MASS J05381330−0251329	SO 401	13.30±0.02	13.16±0.02	13.11±0.04	13.20±0.05	2004 Oct 9
IRAC J05381386−0235095	SO 406	14.24±0.02	13.65±0.02	12.89±0.04	12.16±0.04	2004 Oct 9
		14.44±0.03	13.75±0.02	12.97±0.03	12.18±0.03	2006 Sep 28
		out	13.71±0.02	out	12.18±0.03	2007 Mar 31
2MASS J05381412−0215597	SO 411,HD 294268	8.38±0.02	8.16±0.02	7.61±0.03	5.80±0.03	2004 Oct 9
2MASS J05381462−0240154	S Ori 47	15.18±0.06	15.26±0.06	14.94±0.10	15.02±0.18	2004 Oct 9
		out	15.14±0.03	out	14.68±0.09	2006 Sep 28
		15.06±0.02	15.12±0.02	14.95±0.04	14.97±0.07	2007 Mar 31/Apr 3
2MASS J05381610−0238049	SO 426,S Ori J053816.0-023805	12.37±0.02	12.27±0.02	12.27±0.03	12.28±0.03	2004 Oct 9
		12.39±0.02	12.26±0.02	12.32±0.03	12.26±0.03	2006 Sep 28
		12.39±0.02	12.27±0.02	12.32±0.03	12.24±0.03	2007 Mar 31
2MASS J05381698−0214463	···	12.98±0.02	out	12.82±0.03	out	2004 Oct 9
		out	12.89±0.02	out	12.86±0.03	2007 Apr 3
2MASS J05381718−0222256	SO 432,S Ori 9	12.33±0.02	12.25±0.02	12.20±0.03	12.18±0.03	2004 Oct 9
		12.33±0.02	12.24±0.02	12.17±0.03	12.20±0.03	2007 Apr 3
2MASS J05381741−0240242	S Ori 27	13.55±0.02	13.43±0.02	13.47±0.04	13.47±0.05	2004 Oct 9
		out	13.46±0.02	out	13.54±0.04	2006 Sep 28
		13.55±0.02	13.42±0.02	13.40±0.03	13.39±0.03	2007 Mar 31/Apr 3
2MASS J05381746−0209236	···	12.05±0.02	out	11.92±0.03	out	2004 Oct 9
2MASS J05381778−0240500	SO 435,S Ori J053817.8-024050	11.75±0.02	11.51±0.02	11.35±0.03	10.71±0.03	2004 Oct 9
		out	11.53±0.02	out	10.71±0.03	2006 Sep 28
		sat	11.50±0.02	11.31±0.03	10.71±0.03	2007 Mar 31/Apr 3
2MASS J05381800−0221103	SO 440	10.82±0.02	10.90±0.02	10.88±0.03	10.80±0.03	2004 Oct 9
		sat	10.89±0.00	10.82±0.03	10.76±0.03	2007 Apr 3
2MASS J05381824−0248143	SO 444	11.63±0.02	11.59±0.02	11.54±0.03	11.32±0.03	2004 Oct 9
		sat	11.63±0.00	11.57±0.03	11.56±0.03	2007 Mar 31/Apr 3
2MASS J05381834−0235385	SO 446,S Ori J053818.2-023539	13.87±0.02	13.78±0.02	13.84±0.05	13.93±0.08	2004 Oct 9
		13.88±0.02	13.78±0.02	13.79±0.05	13.71±0.06	2006 Sep 28
		out	13.78±0.02	out	13.70±0.06	2007 Mar 31
2MASS J05381886−0251388	SO 451	11.13±0.02	10.80±0.02	10.36±0.03	9.73±0.03	2004 Oct 9
2MASS J05381975−0236391	SO 457	sat	sat	4.24±0.03	sat	2004 Oct 9
2MASS J05382021−0238016	SO 460,S Ori J053820.2-023802	11.31±0.02	11.26±0.02	11.21±0.03	11.25±0.03	2004 Oct 9
		sat	11.26±0.02	11.27±0.03	11.17±0.04	2006 Sep 28
		sat	11.23±0.02	11.26±0.03	11.20±0.03	2007 Mar 31
2MASS J05382050−0234089	SO 462,S Ori J053820.5-023409	11.03±0.02	10.64±0.02	10.14±0.03	9.00±0.03	2004 Oct 9
		sat	sat	10.06±0.03	8.91±0.03	2006 Sep 28
		out	sat	out	8.90±0.03	2007 Mar 31
2MASS J05382088−0246132	SO 465,S Ori 31	13.83±0.02	13.67±0.02	13.55±0.04	13.69±0.06	2004 Oct 9
		13.78±0.02	13.63±0.02	13.69±0.03	13.62±0.04	2007 Mar 31/Apr 3
2MASS J05382089−0251280	SO 466	13.47±0.02	13.34±0.02	13.28±0.04	13.43±0.06	2004 Oct 9
2MASS J05382119−0254110	SO 467	11.27±0.02	10.98±0.02	10.71±0.03	10.03±0.03	2004 Oct 9
		sat	out	10.58±0.03	out	2007 Apr 3
2MASS J05382138−0233363	SO 469,S Ori J053821.3-023336	14.00±0.02	13.87±0.02	13.80±0.06	13.82±0.07	2004 Oct 9
		out	13.83±0.02	out	13.68±0.04	2007 Mar 31
2MASS J05382178−0221494	SO 475	10.09±0.02	10.18±0.02	10.14±0.03	10.11±0.03	2004 Oct 9
		sat	sat	10.10±0.03	10.10±0.03	2007 Apr 3
2MASS J05382307−0236493	SO 482	12.21±0.02	11.77±0.02	11.30±0.04	10.80±0.03	2004 Oct 9
2MASS J05382332−0244142	SO 484,S Ori J053823.3-024414	12.34±0.02	12.26±0.02	12.28±0.03	12.32±0.04	2004 Oct 9
		out	12.23±0.02	out	12.19±0.03	2006 Sep 28
		12.25±0.02	12.27±0.02	12.20±0.03	12.25±0.03	2007 Mar 31/Apr 3
2MASS J05382333−0225345	SO 485	11.91±0.02	11.58±0.02	11.23±0.03	10.54±0.03	2004 Oct 9
		sat	out	11.16±0.03	out	2007 Apr 3
2MASS J05382354−0241317	SO 489,S Ori J053823.6-024132	12.18±0.02	12.05±0.02	11.99±0.03	12.04±0.03	2004 Oct 9
		out	12.05±0.02	out	12.03±0.03	2006 Sep 28
		12.11±0.02	12.04±0.02	11.96±0.03	12.03±0.03	2007 Mar 31/Apr 3
2MASS J05382358−0220475	SO 490	11.98±0.02	11.64±0.02	11.25±0.03	10.42±0.03	2004 Oct 9
		11.98±0.02	11.65±0.02	11.29±0.03	10.36±0.03	2007 Apr 3
IRAC J05382512−0248026	S Ori 53	15.91±0.03	15.76±0.03	···	···	2004 Oct 9

TABLE 1 — *Continued*

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
		15.96±0.02	15.85±0.03	15.87±0.07	16.20±0.29	2007 Mar 31/Apr 3
2MASS J05382543−0242412	SO 500,S Ori J053825.4-024241	12.73±0.02	12.36±0.02	12.00±0.03	11.38±0.03	2004 Oct 9
		12.75±0.02	12.31±0.02	11.98±0.03	11.44±0.03	2007 Mar 31/Apr 3
2MASS J05382549−0210228	···	10.79±0.02	out	10.83±0.03	out	2004 Oct 9
2MASS J05382557−0248370	S Ori 45	15.03±0.02	15.05±0.03	14.94±0.10	···	2004 Oct 9
		15.17±0.06	15.10±0.06	14.98±0.07	14.85±0.09	2007 Mar 31/Apr 3
2MASS J05382568−0231216	SO 502,S Ori 18	13.39±0.02	13.35±0.02	13.30±0.04	13.32±0.06	2004 Oct 9
		13.41±0.02	out	13.23±0.03	out	2007 Apr 3
IRAC J05382603−0223049	S Ori 65	17.05±0.07	16.67±0.08	···	···	2004 Oct 9
		17.07±0.06	16.83±0.06	17.19±0.21	16.12±0.40	2007 Apr 3
2MASS J05382623−0240413	SO 509,S Ori J053826.1-024041	13.54±0.02	13.44±0.02	13.46±0.04	13.45±0.06	2004 Oct 9
		13.48±0.02	13.41±0.02	13.33±0.03	13.37±0.04	2007 Mar 31/Apr 3
2MASS J05382684−0238460	SO 514,S Ori J053826.8-023846	12.57±0.02	12.26±0.02	11.99±0.03	11.29±0.03	2004 Oct 9
		out	12.23±0.02	out	11.32±0.03	2007 Apr 3
2MASS J05382725−0245096	SO 518	8.76±0.02	8.35±0.02	8.00±0.03	7.19±0.03	2004 Oct 9
2MASS J05382750−0235041	SO 520,S Ori J053827.4-023504	11.42±0.02	11.13±0.02	10.68±0.03	9.76±0.03	2004 Oct 9
2MASS J05382752−0243325	SO 521,HD 294273	10.07±0.02	10.03±0.02	10.05±0.03	10.07±0.03	2004 Oct 9
		out	sat	out	10.09±0.03	2007 Apr 3
2MASS J05382774−0243009	SO 525	11.10±0.02	11.03±0.02	11.01±0.03	11.03±0.03	2004 Oct 9
		out	11.07±0.02	out	11.05±0.03	2007 Apr 3
2MASS J05382896−0248473	SO 537,S Ori J053829.0-024847	13.09±0.02	12.80±0.02	12.49±0.03	11.94±0.03	2004 Oct 9
		13.18±0.02	12.82±0.02	12.51±0.03	11.93±0.03	2007 Mar 31/Apr 3
2MASS J05382911−0236026	SO 539	11.53±0.02	11.53±0.02	11.47±0.03	11.48±0.03	2004 Oct 9
2MASS J05382915−0216156	SO 540	10.50±0.02	10.40±0.02	10.26±0.03	9.44±0.03	2004 Oct 9
		out	sat	out	9.40±0.03	2007 Apr 3
2MASS J05382961−0225141	SO 545,S Ori 29	13.56±0.02	13.47±0.02	13.57±0.04	13.59±0.06	2004 Oct 9
2MASS J05383008−0221198	SO 550	11.27±0.02	11.33±0.02	11.32±0.03	11.29±0.03	2004 Oct 9
		sat	11.33±0.00	11.31±0.03	11.26±0.03	2007 Apr 3
2MASS J05383098−0234038	SO 557	13.72±0.02	13.63±0.02	13.67±0.04	13.52±0.05	2004 Oct 9
		13.72±0.02	out	13.62±0.03	out	2007 Apr 3
2MASS J05383138−0255032	SO 561,HD 294279	9.63±0.02	9.62±0.02	9.65±0.03	9.64±0.03	2004 Oct 9
		sat	out	9.61±0.03	out	2007 Apr 3
2MASS J05383141−0236338	SO 562	10.30±0.02	9.88±0.02	9.51±0.03	8.51±0.03	2004 Oct 9
		sat	out	9.57±0.03	out	2007 Apr 3
2MASS J05383157−0235148	SO 563	9.74±0.02	9.31±0.02	8.90±0.03	8.32±0.03	2004 Oct 9
		sat	out	8.94±0.03	out	2007 Apr 3
2MASS J05383160−0251268	SO 564	10.80±0.02	10.90±0.02	10.80±0.03	10.82±0.03	2004 Oct 9
IRAC J05383213−0232433	SO 566	11.98±0.02	11.83±0.02	11.51±0.03	10.54±0.03	2004 Oct 9
		11.96±0.02	out	11.65±0.03	out	2007 Apr 3
2MASS J05383244−0229573	SO 568,S Ori 39	14.10±0.02	13.94±0.02	13.89±0.05	13.89±0.06	2004 Oct 9
		14.07±0.02	13.96±0.02	13.93±0.03	14.00±0.05	2007 Apr 3
2MASS J05383284−0235392	SO 572	10.59±0.02	10.52±0.02	10.57±0.03	10.55±0.03	2004 Oct 9
		sat	out	10.50±0.03	out	2007 Apr 3
2MASS J05383302−0239279	SO 576	13.41±0.02	13.35±0.02	13.25±0.04	13.27±0.05	2004 Oct 9
		out	13.26±0.02	out	13.19±0.04	2007 Apr 3
2MASS J05383318−0221038	S Ori 54	14.57±0.10	14.77±0.12	14.59±0.14	···	2004 Oct 9
		14.82±0.12	14.78±0.12	14.77±0.13	14.50±0.40	2007 Apr 3
2MASS J05383335−0236176	SO 582	10.93±0.02	10.94±0.02	10.93±0.03	10.91±0.03	2004 Oct 9
		sat	out	10.86±0.03	out	2007 Apr 3
2MASS J05383368−0244141	SO 583	7.68±0.02	7.43±0.02	7.03±0.03	6.10±0.03	2004 Oct 9
2MASS J05383388−0245078	SO 585,S Ori J053833.9-024508	12.86±0.02	12.60±0.02	12.33±0.03	11.51±0.03	2004 Oct 9
		12.85±0.02	12.52±0.02	12.32±0.03	11.57±0.03	2007 Apr 3
2MASS J05383405−0236375	SO 587	10.72±0.02	10.55±0.02	10.43±0.03	9.88±0.03	2004 Oct 9
		sat	out	10.37±0.03	out	2007 Apr 3
2MASS J05383422−0234160	SO 590	8.82±0.02	8.78±0.02	8.76±0.03	8.82±0.03	2004 Oct 9
		sat	out	8.82±0.03	out	2007 Apr 3
2MASS J05383431−0235000	SO 592	10.23±0.02	10.29±0.02	10.23±0.03	10.24±0.03	2004 Oct 9
		sat	out	10.22±0.03	out	2007 Apr 3
2MASS J05383460−0241087	SO 598,S Ori J053834.5-024109	11.78±0.02	11.60±0.02	11.48±0.03	10.91±0.03	2004 Oct 9
2MASS J05383479−0234158	SO 601	8.34±0.02	8.39±0.02	8.33±0.03	8.39±0.03	2004 Oct 9
2MASS J05383484−0232521	SO 602	9.13±0.02	9.18±0.02	9.08±0.03	9.11±0.03	2004 Oct 9
		sat	out	9.19±0.03	out	2007 Apr 3
2MASS J05383528−0233131	SO 606	12.73±0.02	12.60±0.02	12.55±0.04	12.56±0.04	2004 Oct 9
		12.73±0.02	out	12.57±0.03	out	2007 Apr 3
2MASS J05383536−0225222	SO 609,S Ori 22	13.34±0.02	13.20±0.02	13.25±0.04	13.23±0.05	2004 Oct 9
		13.29±0.02	13.17±0.02	13.23±0.03	13.22±0.04	2007 Apr 3
2MASS J05383546−0231516	SO 611	10.31±0.02	10.33±0.02	10.29±0.03	10.25±0.03	2004 Oct 9
2MASS J05383587−0230433	SO 616	10.32±0.02	10.30±0.02	10.24±0.03	10.23±0.03	2004 Oct 9
		sat	sat	10.15±0.03	10.23±0.03	2007 Apr 3
2MASS J05383587−0243512	SO 615	8.59±0.02	8.24±0.02	7.98±0.03	7.75±0.03	2004 Oct 9
2MASS J05383654−0233127	SO 620,HD 294271	8.21±0.02	8.23±0.02	8.20±0.03	8.28±0.03	2004 Oct 9
2MASS J05383669−0244136	SO 621,S Ori J053836.7-024414	11.37±0.02	11.29±0.02	11.27±0.03	11.25±0.03	2004 Oct 9
2MASS J05383687−0236432	SO 624	11.87±0.02	11.76±0.02	11.74±0.03	11.76±0.03	2004 Oct 9
		11.86±0.02	out	11.71±0.03	out	2007 Apr 3
2MASS J05383745−0250236	SO 628	11.61±0.02	11.52±0.02	11.56±0.03	11.54±0.03	2004 Oct 9
		sat	out	11.44±0.03	11.53±0.03	2007 Apr 3

TABLE 1 — *Continued*

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
2MASS J05383787−0220397	SO 631	14.13±0.02	14.02±0.02	14.04±0.05	14.41±0.09	2004 Oct 9
		14.16±0.02	14.02±0.02	14.01±0.03	14.18±0.04	2007 Apr 3
2MASS J05383822−0236384	SO 637	10.21±0.02	10.14±0.02	10.16±0.03	10.13±0.03	2004 Oct 9
		sat	out	10.06±0.03	out	2007 Apr 3
2MASS J05383848−0234550	SO 638	8.98±0.02	9.06±0.02	8.86±0.03	8.77±0.03	2004 Oct 9
		sat	out	8.85±0.03	out	2007 Apr 3
2MASS J05383858−0241558	SO 641,S Ori J053838.6-024157	13.30±0.02	13.25±0.02	13.15±0.03	13.25±0.05	2004 Oct 9
2MASS J05383888−0228016	SO 644	14.10±0.02	14.02±0.02	14.02±0.04	14.01±0.09	2004 Oct 9
		14.13±0.02	14.03±0.02	14.03±0.03	14.10±0.07	2007 Apr 3
2MASS J05383902−0245321	SO 646	11.18±0.02	10.84±0.02	10.48±0.03	9.93±0.03	2004 Oct 9
IRAC J05383916−0228052	S Ori 68	18.09±0.23	18.27±0.32	···	···	2007 Apr 3
2MASS J05383922−0253084	SO 648	11.81±0.02	11.80±0.02	11.76±0.03	11.76±0.03	2004 Oct 9
		11.78±0.02	out	11.75±0.03	out	2007 Apr 3
2MASS J05383972−0218376	SO 654	9.01±0.02	9.12±0.02	9.06±0.03	8.98±0.03	2004 Oct 9
		out	sat	out	8.98±0.03	2007 Apr 3
2MASS J05383972−0240197	SO 655	12.54±0.02	12.46±0.02	12.50±0.03	12.50±0.04	2004 Oct 9
2MASS J05383976−0232203	SO 657	13.15±0.02	12.80±0.02	12.45±0.03	11.71±0.03	2004 Oct 9
		13.23±0.02	out	12.46±0.03	out	2007 Apr 3
2MASS J05383981−0256462	···	out	9.32±0.02	out	7.91±0.03	2004 Oct 9
		sat	out	8.79±0.03	out	2007 Apr 3
2MASS J05384008−0250370	SO 658	12.52±0.02	12.46±0.02	12.46±0.03	12.41±0.04	2004 Oct 9
2MASS J05384027−0230185	SO 662	9.81±0.02	9.41±0.02	8.89±0.03	8.09±0.03	2004 Oct 9
		sat	sat	8.84±0.03	8.07±0.03	2007 Apr 3
2MASS J05384053−0233275	SO 663	11.47±0.02	11.23±0.02	10.94±0.03	10.37±0.03	2004 Oct 9
		sat	out	11.01±0.03	out	2007 Apr 3
2MASS J05384129−0237225	SO 669	10.42±0.02	10.40±0.02	10.36±0.03	10.36±0.03	2004 Oct 9
		sat	out	10.29±0.03	out	2007 Apr 3
2MASS J05384135−0236444	SO 670	11.89±0.02	11.84±0.02	11.82±0.03	11.86±0.03	2004 Oct 9
		11.89±0.02	out	11.75±0.03	out	2007 Apr 3
2MASS J05384143−0217016	SO 671	12.20±0.02	12.21±0.02	12.14±0.03	12.13±0.03	2004 Oct 9
2MASS J05384146−0235523	SO 673	12.35±0.02	12.07±0.02	11.48±0.04	10.89±0.03	2004 Oct 9
		12.32±0.02	out	11.66±0.03	out	2007 Apr 3
2MASS J05384159−0230289	SO 674	11.66±0.02	11.53±0.02	11.31±0.03	10.57±0.03	2004 Oct 9
		11.72±0.02	11.59±0.02	11.32±0.03	10.60±0.03	2007 Apr 3
2MASS J05384227−0237147	SO 682	10.41±0.02	10.22±0.02	9.83±0.03	9.20±0.03	2004 Oct 9
		sat	out	9.78±0.03	out	2007 Apr 3
2MASS J05384239−0236044	SO 683	12.56±0.02	12.27±0.02	11.77±0.04	10.84±0.04	2004 Oct 9
		12.63±0.02	out	11.26±0.04	out	2007 Apr 3
2MASS J05384301−0236145	SO 687	10.06±0.02	9.86±0.02	9.59±0.03	8.91±0.03	2004 Oct 9
		sat	out	9.55±0.03	out	2007 Apr 3
2MASS J05384333−0232008	SO 689	11.06±0.02	10.97±0.02	10.96±0.03	10.96±0.03	2004 Oct 9
		sat	out	11.02±0.03	out	2007 Apr 3
2MASS J05384355−0233253	SO 691	10.53±0.02	10.52±0.02	10.44±0.03	10.47±0.03	2004 Oct 9
		sat	out	10.49±0.03	out	2007 Apr 3
2MASS J05384375−0252427	SO 692	11.80±0.02	11.73±0.02	11.67±0.03	11.69±0.03	2004 Oct 9
2MASS J05384386−0237068	SO 694	11.24±0.02	10.89±0.02	10.60±0.03	9.84±0.03	2004 Oct 9
		sat	out	10.76±0.03	out	2007 Apr 3
2MASS J05384411−0236062	σ Ori C	9.05±0.03	8.96±0.03	9.00±0.03	8.72±0.04	2004 Oct 9
		sat	out	8.92±0.03	out	2007 Apr 3
2MASS J05384423−0240197	SO 697	9.79±0.02	9.41±0.02	8.90±0.03	7.84±0.03	2004 Oct 9
2MASS J05384424−0232336	SO 696	9.89±0.02	9.86±0.02	9.80±0.03	9.79±0.03	2004 Oct 9
		sat	out	9.86±0.03	out	2007 Apr 3
2MASS J05384448−0240376	SO 700,S Ori J053844.4-024037	13.32±0.02	13.27±0.03	13.07±0.04	12.95±0.08	2004 Oct 9
2MASS J05384449−0240305	SO 701,S Ori J053844.4-024030	12.24±0.02	12.21±0.02	12.24±0.04	12.25±0.05	2004 Oct 9
IRAC J05384458−0255136	S Ori J053844.5-025512	17.11±0.06	16.95±0.09	···	···	2004 Oct 9
2MASS J05384480−0233576	SO 706	9.20±0.02	9.25±0.02	9.17±0.03	9.15±0.03	2004 Oct 9
		sat	out	9.21±0.03	out	2007 Apr 3
2MASS J05384527−0237292	SO 707	11.43±0.02	11.12±0.02	10.79±0.03	10.34±0.03	2004 Oct 9
		sat	out	10.51±0.03	out	2007 Apr 3
2MASS J05384537−0241594	SO 710	10.52±0.02	10.26±0.02	9.96±0.03	8.94±0.03	2004 Oct 9
2MASS J05384561−0235588	σ Ori D	7.37±0.02	7.38±0.02	7.40±0.03	7.20±0.03	2004 Oct 9
2MASS J05384597−0245231	SO 714,S Ori J053845.9-024523	12.35±0.02	12.27±0.02	12.24±0.03	12.27±0.04	2004 Oct 9
2MASS J05384684−0236435	SO 721	12.01±0.02	11.95±0.02	11.90±0.03	11.94±0.03	2004 Oct 9
		12.03±0.02	out	11.88±0.03	out	2007 Apr 3
2MASS J05384714−0257557	S Ori J053847.2-025756	out	12.62±0.02	out	11.56±0.03	2004 Oct 9
2MASS J05384718−0234368	SO 723	10.49±0.02	10.16±0.02	9.98±0.03	9.33±0.03	2004 Oct 9
		sat	out	9.79±0.03	out	2007 Apr 3
2MASS J05384719−0235405	SO 724,HD 37479	6.92±0.02	6.83±0.02	6.77±0.03	6.63±0.03	2004 Oct 9
2MASS J05384746−0235252	SO 726	10.19±0.02	9.92±0.02	9.56±0.03	8.68±0.03	2004 Oct 9
		sat	out	8.91±0.03	out	2007 Apr 3
2MASS J05384754−0227119	SO 728,S Ori J053847.5-022711	10.94±0.02	10.85±0.02	10.84±0.04	10.69±0.05	2004 Oct 9
		sat	10.82±0.01	10.72±0.03	10.57±0.03	2007 Mar 30/Apr 3
2MASS J05384766−0230373	SO 730,S Ori 6	12.23±0.02	12.16±0.02	12.10±0.03	12.18±0.04	2004 Oct 9
		12.21±0.02	12.12±0.02	12.07±0.03	12.11±0.03	2007 Mar 30/Apr 3
2MASS J05384791−0237192	SO 733	9.84±0.02	9.43±0.02	8.94±0.03	8.01±0.03	2004 Oct 9
		sat	out	9.19±0.03	out	2007 Apr 3

TABLE 1 — *Continued*

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
2MASS J05384803−0227141	SO 736	8.62±0.02	8.33±0.02	8.03±0.03	7.51±0.03	2004 Oct 9
2MASS J05384809−0228536	SO 738,S Ori 15	12.98±0.02	12.70±0.02	12.37±0.03	11.56±0.03	2004 Oct 9
		13.01±0.02	12.70±0.02	12.37±0.03	11.57±0.03	2007 Mar 30/Apr 3
2MASS J05384818−0244007	SO 739,S Ori J053848.1-024401	12.57±0.02	12.26±0.02	11.93±0.03	11.21±0.03	2004 Oct 9
2MASS J05384828−0236409	SO 740	10.85±0.02	10.81±0.02	10.82±0.03	10.78±0.03	2004 Oct 9
		sat	out	10.71±0.03	out	2007 Apr 3
2MASS J05384868−0236162	SO 742	11.00±0.02	11.03±0.02	11.02±0.03	10.99±0.03	2004 Oct 9
		sat	out	10.91±0.03	out	2007 Apr 3
2MASS J05384917−0238222	SO 747	10.33±0.02	10.28±0.02	10.21±0.03	10.25±0.03	2004 Oct 9
2MASS J05384921−0241251	SO 748	10.57±0.02	10.54±0.02	10.54±0.03	10.52±0.03	2004 Oct 9
2MASS J05384928−0223575	SO 750,S Ori J053849.2-022358	12.79±0.02	12.38±0.02	12.02±0.03	11.60±0.03	2004 Oct 9
		out	12.44±0.02	out	11.69±0.03	2007 Mar 30
IRAC J05384959−0249332	S Ori J053849.5-024934	15.41±0.02	15.11±0.03	14.85±0.08	14.11±0.09	2004 Oct 9
2MASS J05384970−0234526	SO 754	11.64±0.02	11.38±0.02	11.15±0.03	10.58±0.03	2004 Oct 9
		11.68±0.02	out	11.15±0.03	out	2007 Mar 30/Apr 3
2MASS J05384993−0241228	SO 757	11.49±0.02	11.57±0.02	11.55±0.03	11.53±0.03	2004 Oct 9
2MASS J05385003−0237354	SO 758	11.82±0.02	11.78±0.02	11.70±0.03	11.77±0.03	2004 Oct 9
		11.84±0.02	out	11.66±0.03	out	2007 Apr 3
2MASS J05385038−0226477	SO 759	11.30±0.02	11.23±0.02	11.09±0.03	10.85±0.03	2004 Oct 9
		sat	11.12±0.00	11.09±0.03	10.85±0.03	2007 Mar 30/Apr 3
2MASS J05385060−0242429	SO 762,S Ori J053850.6-024244	12.38±0.02	12.09±0.02	11.70±0.03	11.05±0.03	2004 Oct 9
2MASS J05385077−0236267	SO 765	11.91±0.02	11.85±0.02	11.86±0.03	11.81±0.03	2004 Oct 9
		11.91±0.02	out	11.74±0.03	out	2007 Mar 30/Apr 3
2MASS J05385100−0249140	SO 767	13.80±0.02	13.72±0.02	13.56±0.04	13.76±0.06	2004 Oct 9
2MASS J05385145−0236205	SO 770	11.24±0.02	11.13±0.02	11.14±0.03	11.00±0.03	2004 Oct 9
		sat	out	11.03±0.03	out	2007 Mar 30/Apr 3
2MASS J05385173−0236033	SO 773	11.70±0.02	11.63±0.02	11.60±0.03	11.60±0.03	2004 Oct 9
		sat	out	11.54±0.03	out	2007 Mar 30/Apr 3
2MASS J05385200−0246436	SO 774	10.17±0.02	9.93±0.02	9.57±0.03	8.45±0.03	2004 Oct 9
2MASS J05385317−0243528	SO 785	11.11±0.02	11.01±0.02	10.98±0.03	11.00±0.03	2004 Oct 9
2MASS J05385337−0233229	SO 787	9.58±0.02	9.53±0.02	9.49±0.03	9.51±0.03	2004 Oct 9
		sat	sat	9.56±0.03	9.59±0.03	2007 Apr 3
2MASS J05385410−0249297	SO 791	10.12±0.02	10.13±0.02	9.98±0.03	10.13±0.03	2004 Oct 9
2MASS J05385434−0240029	SO 795	13.07±0.02	13.01±0.02	12.94±0.04	12.96±0.04	2004 Oct 9
2MASS J05385492−0228583	SO 797	12.62±0.02	12.53±0.02	12.47±0.03	12.53±0.04	2004 Oct 9
		12.61±0.02	12.52±0.02	12.52±0.03	12.56±0.04	2007 Mar 30/Apr 3
2MASS J05385492−0240337	SO 798,S Ori J053854.9-024034	13.73±0.02	13.43±0.02	12.97±0.04	12.55±0.04	2004 Oct 9
2MASS J05385542−0241208	SO 802,S Ori J053855.4-024121	12.95±0.02	12.54±0.02	12.15±0.03	11.30±0.03	2004 Oct 9
		13.03±0.02	out	12.30±0.03	out	2007 Apr 3
2MASS J05385831−0216101	SO 818	10.46±0.02	10.37±0.02	10.17±0.03	10.09±0.03	2004 Oct 9
		out	sat	out	10.28±0.03	2007 Apr 3
IRAC J05385857−0252265	S Ori J053858.6-025228	16.22±0.03	15.89±0.04	···	···	2004 Oct 9
2MASS J05385911−0247133	SO 823	9.13±0.02	8.55±0.02	8.22±0.03	7.94±0.03	2004 Oct 9
2MASS J05385922−0233514	SO 827	10.89±0.02	10.45±0.02	10.10±0.03	9.29±0.03	2004 Oct 9
		sat	out	9.97±0.03	out	2007 Mar 30/Apr 3
2MASS J05385955−0245080	···	10.83±0.02	10.84±0.02	10.85±0.03	10.85±0.03	2004 Oct 9
		sat	out	10.78±0.03	out	2007 Apr 3
IRAC J05390030−0237060	S Ori 71	15.23±0.02	14.95±0.03	14.67±0.08	13.79±0.07	2004 Oct 9
		15.15±0.02	14.79±0.02	14.47±0.03	13.71±0.04	2007 Mar 30/Apr 3
IRAC J05390079−0221418	S Ori 56	16.09±0.03	15.82±0.03	···	···	2004 Oct 9
		16.15±0.02	15.81±0.02	15.47±0.06	14.95±0.10	2007 Mar 30/Apr 3
2MASS J05390115−0236388	SO 841	12.46±0.02	12.35±0.02	12.45±0.03	12.31±0.03	2004 Oct 9
		12.39±0.02	12.32±0.02	12.31±0.03	12.33±0.03	2007 Mar 30/Apr 3
2MASS J05390136−0218274	SO 844	8.93±0.02	8.52±0.02	8.20±0.03	7.55±0.03	2004 Oct 9
2MASS J05390149−0238564	SO 847,HD 37525	8.07±0.02	8.06±0.02	8.04±0.03	8.07±0.03	2004 Oct 9
		sat	sat	sat	8.14±0.03	2007 Mar 30/Apr 3
2MASS J05390193−0235029	SO 848,S Ori J053902.1-023501	11.53±0.02	11.06±0.02	10.60±0.03	9.85±0.03	2004 Oct 9
		sat	10.98±0.02	10.58±0.03	9.88±0.03	2007 Mar 30/Apr 3
2MASS J05390276−0229558	SO 855	11.47±0.02	11.38±0.02	11.36±0.03	11.38±0.03	2004 Oct 9
		sat	11.43±0.00	11.39±0.03	11.41±0.03	2007 Mar 30/Apr 3
2MASS J05390297−0241272	SO 859	10.31±0.02	9.94±0.02	9.70±0.03	9.14±0.03	2004 Oct 9
		sat	out	9.81±0.03	out	2007 Apr 3
IRAC J05390321−0230200	S Ori 51	15.41±0.02	15.24±0.03	15.10±0.11	···	2004 Oct 9
		15.40±0.02	15.29±0.02	15.19±0.04	15.26±0.07	2007 Mar 30/Apr 3
2MASS J05390357−0246269	SO 865	11.49±0.02	11.22±0.02	10.90±0.03	10.15±0.03	2004 Oct 9
		sat	out	10.81±0.03	out	2007 Apr 3
IRAC J05390360−0225366	S Ori 58	16.48±0.05	16.22±0.05	···	···	2004 Oct 9
		16.48±0.03	16.19±0.03	15.78±0.06	15.73±0.20	2007 Mar 30/Apr 3
2MASS J05390387−0220081	SO 866	12.42±0.02	12.06±0.02	11.66±0.03	10.96±0.03	2004 Oct 9
		12.41±0.02	12.05±0.02	11.73±0.03	10.94±0.03	2007 Mar 30/Apr 3
2MASS J05390443−0252309	SO 869	10.69±0.02	10.69±0.02	10.62±0.03	10.63±0.03	2004 Oct 9
2MASS J05390449−0238353	SO 870,S Ori 17	13.45±0.02	13.34±0.02	13.26±0.04	13.21±0.05	2004 Oct 9
		13.45±0.02	13.36±0.02	13.27±0.03	13.34±0.03	2007 Mar 30/Apr 3
2MASS J05390458−0241493	SO 871	10.99±0.02	10.65±0.02	10.27±0.03	9.59±0.03	2004 Oct 9
		sat	out	10.32±0.03	out	2007 Apr 3
2MASS J05390524−0233005	SO 877	12.21±0.02	12.13±0.02	12.12±0.03	12.16±0.03	2004 Oct 9

TABLE 1 — *Continued*

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
		12.21±0.02	12.17±0.02	12.13±0.03	12.19±0.03	2007 Apr 3
2MASS J05390540−0232303	SO 879	10.57±0.02	10.63±0.02	10.59±0.03	10.56±0.03	2004 Oct 9
		sat	sat	10.53±0.03	10.55±0.03	2007 Mar 30/Apr 3
2MASS J05390756−0212145	· · ·	13.07±0.02	out	12.47±0.03	out	2004 Oct 9
		out	12.73±0.02	out	11.84±0.03	2007 Apr 3
2MASS J05390759−0228234	SO 896	11.73±0.02	11.70±0.02	11.67±0.03	11.69±0.03	2004 Oct 9
		11.76±0.02	11.69±0.02	11.68±0.03	11.68±0.03	2007 Mar 30/Apr 3
2MASS J05390760−0232391	SO 897	9.77±0.02	9.63±0.02	9.59±0.03	9.34±0.03	2004 Oct 9
		sat	sat	9.68±0.03	9.40±0.03	2007 Apr 3
2MASS J05390758−0229055	SO 895,S Ori 20	13.56±0.02	13.53±0.02	13.35±0.04	13.62±0.07	2004 Oct 9
		13.60±0.02	13.52±0.02	13.48±0.03	13.52±0.04	2007 Mar 30/Apr 3
2MASS J05390808−0228447	SO 901,S Ori 8	12.91±0.02	12.78±0.02	12.61±0.03	12.56±0.04	2004 Oct 9
		12.79±0.02	12.56±0.02	12.31±0.03	12.09±0.03	2007 Mar 30/Apr 3
2MASS J05390821−0232284	SO 902,S Ori 7	12.59±0.02	12.52±0.02	12.58±0.03	12.67±0.04	2004 Oct 9
		12.61±0.02	12.52±0.02	12.48±0.03	12.51±0.03	2007 Apr 3
2MASS J05390853−0251465	SO 905	10.82±0.02	10.73±0.02	10.65±0.03	10.37±0.03	2004 Oct 9
2MASS J05390878−0231115	SO 908	11.05±0.02	10.88±0.02	10.82±0.03	10.38±0.03	2004 Oct 9
		sat	10.94±0.00	10.78±0.03	10.35±0.03	2007 Mar 30/Apr 3
2MASS J05390893−0257049	· · ·	out	12.05±0.02	· · ·	12.01±0.03	2004 Oct 9
2MASS J05390894−0239579	SO 911,S Ori 25	13.28±0.05	13.21±0.06	13.23±0.07	13.01±0.07	2004 Oct 9
		13.33±0.06	13.18±0.05	13.05±0.06	13.20±0.07	2007 Apr 3
2MASS J05390921−0256347	SO 913,HD 37545	9.33±0.02	9.34±0.02	9.30±0.03	9.29±0.03	2004 Oct 9
2MASS J05390928−0219130	SO 914	11.57±0.02	11.51±0.02	11.49±0.03	11.52±0.03	2004 Oct 9
		sat	11.52±0.00	11.52±0.03	11.51±0.03	2007 Apr 3
2MASS J05391001−0228116	SO 917,S Ori J053909.9-022814	13.40±0.02	13.38±0.02	13.38±0.04	13.22±0.06	2004 Oct 9
		13.43±0.02	13.31±0.02	13.32±0.03	13.35±0.04	2007 Apr 3
2MASS J05391003−0242425	SO 918	11.79±0.02	11.89±0.02	11.84±0.03	11.83±0.03	2004 Oct 9
		11.83±0.02	out	11.86±0.03	out	2007 Apr 3
IRAC J05391081−0237146	S Ori 50	15.64±0.03	15.33±0.03	· · ·	· · ·	2004 Oct 9
		15.67±0.02	15.57±0.02	15.37±0.04	16.56±0.41	2007 Mar 30/Apr 3
2MASS J05391120−0225490	SO 924	10.39±0.02	10.41±0.02	10.36±0.03	10.40±0.03	2004 Oct 9
		sat	sat	10.36±0.03	10.38±0.03	2007 Apr 3
2MASS J05391139−0233327	SO 925,S Ori J053911.4-023333	13.22±0.02	13.13±0.02	13.03±0.03	13.01±0.04	2004 Oct 9
		13.21±0.02	out	13.03±0.03	out	2007 Mar 30/Apr 3
2MASS J05391151−0231065	SO 927	9.65±0.02	9.06±0.02	8.35±0.03	7.21±0.03	2004 Oct 9
		sat	sat	9.51±0.03	8.48±0.03	2007 Apr 3
2MASS J05391163−0236028	SO 929	10.61±0.02	10.58±0.02	10.57±0.03	10.54±0.03	2004 Oct 9
		sat	sat	10.51±0.03	10.48±0.03	2007 Mar 30/Apr 3
2MASS J05391183−0227409	SO 931,S Ori 1	12.52±0.02	12.39±0.02	12.40±0.03	12.31±0.04	2004 Oct 9
		12.41±0.02	12.32±0.02	12.32±0.03	12.31±0.03	2007 Apr 3
2MASS J05391232−0230064	SO 933	11.37±0.02	11.28±0.02	11.23±0.03	11.25±0.03	2004 Oct 9
		sat	11.27±0.00	11.22±0.03	11.28±0.03	2007 Apr 3
2MASS J05391289−0224537	S Ori J053912.8-022453	15.04±0.02	14.94±0.03	14.73±0.08	· · ·	2004 Oct 9
		15.06±0.02	14.90±0.02	14.73±0.04	14.35±0.08	2007 Apr 3
2MASS J05391308−0237509	SO 936,S Ori 30	13.84±0.02	13.53±0.02	13.32±0.04	12.62±0.04	2004 Oct 9
		13.79±0.02	13.49±0.02	13.21±0.03	12.63±0.03	2007 Mar 30/Apr 3
2MASS J05391346−0237391	SO 940	12.21±0.02	12.10±0.02	12.12±0.03	12.07±0.03	2004 Oct 9
		12.20±0.02	12.09±0.02	12.06±0.03	12.11±0.03	2007 Mar 30/Apr 3
2MASS J05391447−0228333	SO 946,S Ori J053914.5-022834	12.16±0.02	12.06±0.02	12.05±0.03	12.06±0.03	2004 Oct 9
		12.16±0.02	12.07±0.02	12.04±0.03	12.00±0.03	2007 Apr 3
2MASS J05391453−0219367	SO 947	11.07±0.02	11.05±0.02	11.02±0.03	11.01±0.03	2004 Oct 9
		out	11.00±0.02	out	11.02±0.03	2007 Apr 3
2MASS J05391483−0244415	SO 951	11.43±0.02	11.46±0.02	11.40±0.03	11.45±0.03	2004 Oct 9
		sat	out	11.37±0.03	out	2007 Apr 3
2MASS J05391506−0231376	SO 956,HD 37564	7.74±0.02	7.63±0.02	7.59±0.03	7.44±0.03	2004 Oct 9
2MASS J05391510−0240475	SO 957,S Ori 16	13.29±0.02	13.22±0.02	13.14±0.04	13.19±0.06	2004 Oct 9
		out	out	13.14±0.03	out	2007 Apr 3
2MASS J05391525−0221507	SO 959,S Ori 38	13.81±0.02	13.47±0.02	13.08±0.04	12.70±0.04	2004 Oct 9
		out	13.39±0.02	out	12.56±0.03	2007 Apr 3
2MASS J05391582−0236507	SO 967	11.55±0.02	11.29±0.02	11.10±0.03	10.68±0.03	2004 Oct 9
		sat	11.15±0.00	10.88±0.03	10.50±0.03	2007 Apr 3
2MASS J05391659−0222242	SO 972	9.75±0.02	9.84±0.02	9.79±0.03	9.76±0.03	2004 Oct 9
		out	sat	out	9.72±0.03	2007 Apr 3
2MASS J05391699−0241171	SO 976	13.09±0.02	13.04±0.02	13.02±0.04	13.02±0.05	2004 Oct 9
		13.09±0.02	out	13.04±0.03	out	2007 Apr 3
2MASS J05391717−0225433	SO 978	11.77±0.02	11.70±0.02	11.64±0.03	11.65±0.03	2004 Oct 9
IRAC J05391803−0228547	S Ori 69	18.53±0.12	· · ·	· · ·	· · ·	2004 Oct 9
		17.98±0.04	17.66±0.05	· · ·	· · ·	2007 Apr 3
2MASS J05391807−0229284	SO 981	10.05±0.02	10.04±0.02	9.99±0.03	10.01±0.03	2004 Oct 9
		sat	sat	10.01±0.03	10.01±0.03	2007 Apr 3
2MASS J05391813−0252563	S Ori J053918.1-025257	14.53±0.02	14.26±0.02	13.83±0.05	13.42±0.06	2004 Oct 9
2MASS J05391883−0230531	SO 984	9.87±0.02	9.55±0.02	9.29±0.03	8.53±0.03	2004 Oct 9
		sat	sat	sat	8.14±0.00	2007 Apr 3
2MASS J05392023−0238258	SO 999,S Ori 5	12.39±0.02	12.29±0.02	12.26±0.03	12.26±0.04	2004 Oct 9
		12.39±0.02	out	12.05±0.03	out	2007 Apr 3
2MASS J05392044−0227368	SO 1000,S Ori J053920.5-022737	10.99±0.02	10.93±0.02	10.90±0.03	10.90±0.03	2004 Oct 9

TABLE 1 — *Continued*

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
		sat	10.96±0.00	10.93±0.03	10.90±0.03	2007 Apr 3
2MASS J05392097−0230334	SO 1005,S Ori 3	12.09±0.02	11.93±0.02	11.92±0.03	11.88±0.03	2004 Oct 9
		12.06±0.02	11.98±0.02	11.89±0.03	11.89±0.03	2007 Apr 3
2MASS J05392174−0244038	SO 1009	10.13±0.02	10.12±0.02	10.07±0.03	10.10±0.03	2004 Oct 9
2MASS J05392224−0245524	SO 1013,S Ori J053922.2-024552	14.11±0.02	14.05±0.02	13.93±0.05	14.11±0.09	2004 Oct 9
2MASS J05392286−0233330	SO 1017	11.66±0.02	11.65±0.02	11.58±0.03	11.62±0.03	2004 Oct 9
		11.75±0.02	out	11.64±0.03	out	2007 Apr 3
2MASS J05392319−0246557	SO 1019,S Ori 28	13.96±0.02	13.87±0.02	13.80±0.05	13.94±0.07	2004 Oct 9
2MASS J05392341−0240575	S Ori 42	14.76±0.02	14.41±0.02	14.24±0.06	13.43±0.06	2004 Oct 9
2MASS J05392435−0234013	SO 1027	11.79±0.02	11.83±0.02	11.76±0.03	11.78±0.03	2004 Oct 9
		11.87±0.02	out	11.78±0.03	out	2007 Apr 3
2MASS J05392456−0220441	SO 1030	10.47±0.02	10.49±0.02	10.37±0.03	10.41±0.03	2004 Oct 9
		out	sat	out	10.37±0.03	2007 Apr 3
2MASS J05392519−0238220	SO 1036	9.13±0.02	8.83±0.02	8.56±0.03	7.97±0.03	2004 Oct 9
		sat	out	8.75±0.03	out	2007 Apr 3
2MASS J05392524−0227479	SO 1037	14.29±0.02	14.25±0.02	14.41±0.06	14.16±0.10	2004 Oct 9
		14.31±0.02	14.20±0.02	14.26±0.03	14.41±0.07	2007 Apr 3
2MASS J05392561−0234042	SO 1043	11.96±0.02	11.85±0.02	11.83±0.03	11.85±0.03	2004 Oct 9
		11.96±0.02	out	11.86±0.03	out	2007 Apr 3
2MASS J05392560−0238436	HH 446	12.88±0.02	12.14±0.02	11.24±0.03	9.81±0.03	2004 Oct 9
2MASS J05392633−0228376	SO 1050,S Ori 2	12.18±0.02	11.94±0.02	11.70±0.03	10.85±0.03	2004 Oct 9
		12.12±0.02	11.86±0.02	11.65±0.03	10.99±0.03	2007 Apr 3
2MASS J05392639−0215034	Haro 5-22	9.27±0.02	out	8.38±0.03	out	2004 Oct 9
2MASS J05392647−0226154	SO 1052	12.30±0.02	12.24±0.02	12.24±0.03	12.18±0.03	2004 Oct 9
		12.30±0.02	12.25±0.02	12.21±0.03	12.17±0.03	2007 Mar 30/Apr 3
2MASS J05392650−0252152	SO 1053	11.84±0.02	11.75±0.02	11.76±0.03	11.75±0.03	2004 Oct 9
2MASS J05392677−0242583	SO 1057	11.85±0.02	11.77±0.02	11.62±0.03	11.18±0.03	2004 Oct 9
2MASS J05392680−0219247	SO 1058	10.99±0.02	10.98±0.02	10.99±0.03	10.96±0.03	2004 Oct 9
2MASS J05392685−0236561	SO 1059,S Ori 36	13.76±0.02	13.49±0.02	13.11±0.04	12.46±0.04	2004 Oct 9
		13.79±0.02	out	13.13±0.03	out	2007 Apr 3
2MASS J05392883−0217513	SO 1071	11.43±0.02	11.44±0.02	11.38±0.03	11.40±0.03	2004 Oct 9
2MASS J05392935−0227209	SO 1075	10.75±0.02	10.20±0.02	9.66±0.03	8.82±0.03	2004 Oct 9
		sat	sat	9.32±0.03	8.50±0.03	2007 Apr 3
IRAC J05392937−0246364	S Ori J053929.4-024636	15.49±0.06	15.54±0.07	···	···	2004 Oct 9
2MASS J05393132−0248528	SO 1083	12.43±0.02	12.36±0.02	12.30±0.03	12.42±0.04	2004 Oct 9
2MASS J05393234−0227571	SO 1092	10.20±0.02	10.27±0.02	10.23±0.03	10.21±0.03	2004 Oct 9
		sat	sat	10.26±0.03	10.13±0.03	2007 Mar 30
IRAC J05393242−0252199	S Ori J053932.4-025220	16.63±0.12	16.62±0.12	···	···	2004 Oct 9
2MASS J05393256−0239440	SO 1094	9.78±0.02	9.75±0.02	9.70±0.03	9.68±0.03	2004 Oct 9
2MASS J05393291−0247492	SO 1097	10.74±0.02	10.74±0.02	10.72±0.03	10.71±0.03	2004 Oct 9
2MASS J05393378−0220398	SO 1104	11.30±0.02	11.32±0.02	11.23±0.03	11.10±0.03	2004 Oct 9
2MASS J05393432−0238468	SO 1108,S Ori 21	13.40±0.02	13.25±0.02	13.29±0.04	13.17±0.05	2004 Oct 9
		13.38±0.02	out	13.19±0.03	out	2007 Mar 30
2MASS J05393511−0247299	SO 1113	10.76±0.02	10.81±0.02	10.80±0.03	10.76±0.03	2004 Oct 9
2MASS J05393654−0242171	SO 1123	7.89±0.02	7.90±0.02	7.89±0.03	7.87±0.03	2004 Oct 9
2MASS J05393685−0250247	SO 1129	8.71±0.02	8.79±0.02	8.69±0.03	8.66±0.03	2004 Oct 9
2MASS J05393729−0226567	SO 1133	10.65±0.02	10.64±0.02	10.61±0.03	10.60±0.03	2004 Oct 9
		out	sat	out	10.62±0.03	2007 Mar 30
IRAC J05393752−0230418	S Ori 60	16.55±0.04	16.44±0.08	···	···	2004 Oct 9
		16.64±0.02	16.38±0.03	15.89±0.07	15.03±0.10	2007 Mar 30
2MASS J05393759−0244304	SO 1135,S Ori 14	12.99±0.02	12.90±0.02	12.87±0.04	12.92±0.04	2004 Oct 9
2MASS J05393778−0228443	SO 1137	10.42±0.02	10.39±0.02	10.39±0.03	10.36±0.03	2004 Oct 9
		sat	sat	10.39±0.03	10.34±0.03	2007 Mar 30
2MASS J05393931−0232252	SO 1151,S Ori 4	12.24±0.02	12.12±0.02	12.14±0.03	12.18±0.04	2004 Oct 9
		12.19±0.02	12.12±0.02	12.03±0.03	12.11±0.03	2007 Mar 30
2MASS J05393938−0217045	SO 1152	9.32±0.02	8.96±0.02	8.61±0.03	8.17±0.03	2004 Oct 9
2MASS J05393982−0231217	SO 1153,HH 444	8.68±0.02	8.06±0.02	7.55±0.03	6.49±0.03	2004 Oct 9
2MASS J05393982−0233159	SO 1154,HH 445	8.80±0.02	8.38±0.02	8.11±0.03	7.55±0.03	2004 Oct 9
2MASS J05393998−0243097	SO 1155	8.85±0.02	8.55±0.02	8.22±0.03	7.33±0.03	2004 Oct 9
2MASS J05394017−0220480	SO 1156	9.93±0.02	9.69±0.02	9.56±0.03	8.95±0.03	2004 Oct 9
2MASS J05394057−0225468	SO 1163,HD 294299	9.99±0.02	9.91±0.02	9.91±0.03	9.96±0.03	2004 Oct 9
		out	sat	out	10.01±0.03	2007 Mar 30
2MASS J05394057−0239123	SO 1162	14.15±0.02	14.01±0.02	14.01±0.05	14.05±0.08	2004 Oct 9
		14.09±0.02	out	13.93±0.03	out	2007 Mar 30
2MASS J05394097−0216243	···	11.05±0.02	out	10.33±0.03	out	2004 Oct 9
IRAC J05394206−0230316	S Ori 62	17.05±0.06	16.80±0.11	···	···	2004 Oct 9
		16.97±0.03	16.78±0.04	16.89±0.16	16.99±0.48	2007 Mar 30
2MASS J05394299−0213333	···	12.71±0.02	out	12.65±0.03	out	2004 Oct 9
2MASS J05394318−0232433	SO 1182,S Ori J053943.2-023243	11.12±0.02	10.79±0.02	10.51±0.03	9.89±0.03	2004 Oct 9
		sat	sat	10.21±0.03	9.79±0.03	2007 Mar 30
2MASS J05394354−0246275	SO 1186	12.31±0.02	12.33±0.02	12.28±0.03	12.29±0.04	2004 Oct 9
2MASS J05394358−0247318	SO 1188,S Ori 32	13.88±0.02	13.91±0.02	13.78±0.05	13.70±0.07	2004 Oct 9
2MASS J05394411−0231092	SO 1189	10.26±0.02	10.28±0.02	10.24±0.03	10.21±0.03	2004 Oct 9
		sat	sat	10.25±0.03	10.23±0.03	2007 Mar 30
2MASS J05394433−0233027	SO 1191,S Ori 11	13.03±0.02	12.97±0.02	12.99±0.04	12.95±0.05	2004 Oct 9
		13.08±0.02	12.99±0.02	12.78±0.03	13.04±0.04	2007 Mar 30

TABLE 1 — *Continued*

ID[a]	Name[b]	[3.6]	[4.5]	[5.8]	[8.0]	Date
2MASS J05394450−0224432	SO 1193,S Ori 10	12.64±0.02	12.37±0.02	12.05±0.03	11.31±0.03	2004 Oct 9
		out	12.35±0.02	out	11.33±0.03	2007 Mar 30
IRAC J05394456−0259589	S Ori J053944.5-025959	out	16.40±0.04	out	···	2004 Oct 9
2MASS J05394619−0240320	SO 1204,HD 37633	8.80±0.02	8.86±0.02	8.81±0.03	8.85±0.03	2004 Oct 9
2MASS J05394645−0224230	S Ori J053946.5-022423	15.22±0.02	14.96±0.03	14.73±0.08	14.23±0.09	2004 Oct 9
		out	15.01±0.02	out	14.14±0.04	2007 Mar 30
2MASS J05394661−0226313	SO 1207,S Ori J053946.6-022631	11.78±0.02	11.65±0.02	11.66±0.03	11.70±0.03	2004 Oct 9
		out	11.65±0.02	out	11.72±0.03	2007 Mar 30
2MASS J05394741−0226162	SO 1215	9.20±0.02	9.23±0.02	9.20±0.03	9.16±0.03	2004 Oct 9
		out	sat	out	9.21±0.03	2007 Mar 30
2MASS J05394770−0236230	SO 1216	12.32±0.02	12.21±0.02	12.20±0.03	12.27±0.03	2004 Oct 9
		12.29±0.02	out	12.21±0.03	out	2007 Mar 30
2MASS J05394784−0232248	SO 1217	10.01±0.02	9.98±0.02	9.94±0.03	9.96±0.03	2004 Oct 9
		sat	sat	9.87±0.03	9.95±0.03	2007 Mar 30
2MASS J05394799−0240320	SO 1219	11.26±0.02	11.29±0.02	11.18±0.03	11.20±0.03	2004 Oct 9
2MASS J05394806−0245571	SO 1220,S Ori J053948.1-024557	11.83±0.02	11.76±0.02	11.71±0.03	11.80±0.03	2004 Oct 9
2MASS J05394891−0229110	SO 1224	11.98±0.02	11.93±0.02	11.93±0.03	11.87±0.03	2004 Oct 9
		12.01±0.02	11.91±0.02	11.93±0.03	11.89±0.03	2007 Mar 30
2MASS J05394943−0217322	SO 1229	10.74±0.02	10.82±0.02	10.74±0.03	10.78±0.03	2004 Oct 9
2MASS J05394944−0223459	SO 1230,S Ori J053949.3-022346	12.09±0.02	11.88±0.02	11.61±0.03	11.02±0.03	2004 Oct 9
IRAC J05394953−0231297	S Ori J053949.5-023130	16.63±0.07	16.52±0.07	···	···	2004 Oct 9
		16.59±0.06	16.43±0.06	16.31±0.14	···	2007 Mar 30
2MASS J05395038−0243307	SO 1235	10.60±0.02	10.57±0.02	10.62±0.03	10.59±0.03	2004 Oct 9
2MASS J05395056−0234137	SO 1238,S Ori J053950.6-023414	12.48±0.02	12.45±0.02	12.38±0.03	12.41±0.04	2004 Oct 9
		12.60±0.02	out	12.43±0.03	out	2007 Mar 30
2MASS J05395118−0222461	SO 1246	12.05±0.02	12.11±0.02	11.99±0.03	11.98±0.04	2004 Oct 9
2MASS J05395172−0222472	SO 1248,S Ori J053951.7-022247	11.20±0.02	10.78±0.02	10.34±0.03	9.58±0.03	2004 Oct 9
2MASS J05395248−0232023	SO 1250	10.55±0.02	10.50±0.02	10.49±0.03	10.48±0.03	2004 Oct 9
		sat	sat	10.40±0.03	10.48±0.03	2007 Mar 30
2MASS J05395253−0243223	SO 1251	9.55±0.02	9.61±0.02	9.55±0.03	9.52±0.03	2004 Oct 9
2MASS J05395313−0243083	SO 1256	10.16±0.02	10.14±0.02	10.17±0.03	10.13±0.03	2004 Oct 9
2MASS J05395362−0233426	SO 1260	10.76±0.02	10.38±0.02	10.11±0.03	9.23±0.03	2004 Oct 9
2MASS J05395421−0227326	SO 1266,S Ori J053954.2-022733	12.18±0.02	11.86±0.02	11.41±0.03	10.04±0.03	2004 Oct 9
2MASS J05395428−0224385	SO 1267	9.60±0.02	9.41±0.02	9.35±0.03	8.81±0.03	2004 Oct 9
2MASS J05395433−0237189	SO 1268,S Ori J053954.3-023719	13.44±0.02	13.29±0.02	13.40±0.04	13.36±0.05	2004 Oct 9
		13.40±0.02	out	13.21±0.03	out	2007 Mar 30
2MASS J05395437−0244476	SO 1269	9.16±0.02	9.22±0.02	9.13±0.03	9.13±0.03	2004 Oct 9
2MASS J05395465−0246341	SO 1274	9.19±0.02	8.88±0.02	8.56±0.03	7.66±0.03	2004 Oct 9
2MASS J05395484−0219299	SO 1275	11.22±0.02	11.22±0.02	11.22±0.03	11.17±0.03	2004 Oct 9
2MASS J05395594−0220366	SO 1282	11.13±0.02	11.12±0.02	11.02±0.03	11.01±0.03	2004 Oct 9
2MASS J05395645−0238034	SO 1285,S Ori J053956.4-023804	11.95±0.02	11.62±0.02	11.28±0.03	10.54±0.03	2004 Oct 9
IRAC J05395682−0253146	S Ori J053956.8-025315	16.47±0.04	16.45±0.10	···	···	2004 Oct 9
2MASS J05395753−0232120	SO 1295,S Ori J053957.5-023212	12.10±0.02	12.03±0.02	12.01±0.03	11.95±0.03	2004 Oct 9
2MASS J05395761−0247359	SO 1296	12.21±0.02	12.23±0.02	12.21±0.03	12.19±0.03	2004 Oct 9
2MASS J05395798−0248006	SO 1299	11.29±0.02	11.29±0.02	11.26±0.03	11.27±0.03	2004 Oct 9
2MASS J05395825−0226188	SO 1301,S Ori J053958.2-022619	11.63±0.02	11.53±0.02	11.54±0.03	11.54±0.03	2004 Oct 9
2MASS J05395930−0222543	SO 1307,HD 294298	8.69±0.02	8.68±0.02	8.65±0.03	8.66±0.03	2004 Oct 9
2MASS J05400014−0251594	SO 1316,S Ori J054000.2-025159	12.80±0.02	12.73±0.02	12.73±0.03	12.65±0.04	2004 Oct 9
2MASS J05400101−0219597	SO 1323	11.94±0.02	11.79±0.02	11.84±0.03	11.59±0.03	2004 Oct 9
2MASS J05400195−0221325	SO 1327,S Ori J054001.8-022133	10.65±0.02	10.36±0.02	10.20±0.03	9.69±0.03	2004 Oct 9
2MASS J05400338−0229014	SO 1337	10.72±0.02	10.72±0.02	10.70±0.03	10.68±0.03	2004 Oct 9
2MASS J05400365−0216461	Haro 5-35	10.25±0.02	out	9.49±0.03	out	2004 Oct 9
2MASS J05400453−0236421	SO 1338,S Ori J054004.5-023642	13.67±0.02	13.37±0.02	12.95±0.04	12.27±0.03	2004 Oct 9
2MASS J05400510−0219591	SO 1343	10.43±0.02	10.40±0.02	10.37±0.03	10.33±0.03	2004 Oct 9
2MASS J05400525−0230522	SO 1344,S Ori J054005.3-023052	12.43±0.02	12.15±0.02	11.78±0.03	11.26±0.03	2004 Oct 9
2MASS J05400530−0245380	SO 1347	bad	9.12±0.02	bad	9.03±0.03	2004 Oct 9
2MASS J05400696−0228300	SO 1352	10.11±0.02	10.12±0.02	10.07±0.03	10.13±0.03	2004 Oct 9
IRAC J05400696−0236049	S Ori J054007.0-023604	15.94±0.04	15.78±0.04	···	···	2004 Oct 9
2MASS J05400708−0232446	SO 1353,S Ori J054007.1-023245	12.24±0.02	12.13±0.02	12.20±0.03	12.15±0.03	2004 Oct 9
2MASS J05400867−0232432	SO 1359	10.59±0.02	10.52±0.02	10.49±0.03	10.49±0.03	2004 Oct 9
2MASS J05400889−0233336	SO 1361,HH 447	9.11±0.02	8.66±0.02	8.33±0.03	7.50±0.03	2004 Oct 9
2MASS J05400933−0225067	SO 1362,S Ori J054009.3-022507	11.51±0.02	11.07±0.02	10.67±0.03	9.89±0.03	2004 Oct 9
2MASS J05401274−0228199	SO 1368	10.11±0.02	10.11±0.02	10.06±0.03	10.06±0.03	2004 Oct 9
2MASS J05401286−0222020	SO 1369	8.73±0.02	8.26±0.02	7.83±0.03	7.16±0.03	2004 Oct 9
2MASS J05401471−0222265	SO 1374	10.98±0.02	11.07±0.02	11.01±0.03	10.98±0.03	2004 Oct 9

NOTE. — Entries of "···", "sat", and "out" indicate measurements that are absent because of non-detection, saturation, and a position outside the field of view of the filter, respectively.

[a] 2MASS identifications are from the 2MASS Point Source Catalog. IRAC identifications are based on J2000 coordinates measured with IRAC.[a] SO identifications are from Hernández et al. (2007a).

TABLE 2
Spitzer Photometry for Candidate Members of σ Ori

Name	[3.6]	[4.5]	[5.8]	[8.0]	Date	[24]	Date
2MASS J05375398−0249545[a]	11.10±0.02	10.70±0.02	10.39±0.03	9.80±0.04	2004 Oct 9	7.24±0.04	2004 Mar 17
2MASS J05383446−0253514[b]	10.50±0.02	9.77±0.02	9.14±0.03	8.51±0.03	2004 Oct 9	5.68±0.04	2004 Mar 17
	sat	out	9.54±0.03	out	2007 Apr 3	···	···
IRAC J05384459−0225433	15.75±0.03	15.35±0.03	14.96±0.09	14.35±0.09	2004 Oct 9	···	···
	15.87±0.03	15.52±0.03	14.93±0.04	14.48±0.06	2007 Mar 30/Apr 3	···	···
IRAC J05384729−0235194	14.16±0.20	13.21±0.12	12.72±0.14	11.22±0.13	2007 Apr 3	···	···
IRAC J05391066−0229238	15.77±0.03	15.15±0.03	14.38±0.07	14.04±0.06	2004 Oct 9	···	···
	16.15±0.02	15.68±0.02	14.89±0.04	14.19±0.06	2007 Apr 3	···	···

Note. — Entries of “sat” and “out” indicate measurements that are absent because of saturation and a position outside the field of view of the filter, respectively.

[a] Identified as a candidate member by Sherry et al. (2004). [b] Identified as a candidate member by Caballero (2008).

TABLE 3
IRAC Classifications for Candidate Brown Dwarfs in σ Ori

Name	C07[a]	Z07[b]	SJ08[c]	this work
S Ori 47[d]	no disk	···	no disk	no disk
S Ori 50	no disk	···	no disk	no disk
S Ori 51	no disk	···	no disk	no disk
S Ori 53	? (too faint)	···	no disk	no disk
S Ori 54	···	disk	no disk	no disk
S Ori 55	···	disk	no disk	disk?
S Ori 56	···	disk	no disk	disk
S Ori 58	···	disk	disk?	disk?
S Ori 60	? (too faint)	disk	disk	disk
S Ori 62	? (too faint)	···	no disk	no disk
S Ori 65	···	···	disk	no disk
S Ori 66	···	···	disk?	disk?
S Ori 67	···	···	? (too faint)	? (too faint)
S Ori 68	···	···	? (too faint)	? (too faint)
S Ori 69	···	···	? (too faint)	? (too faint)
S Ori 70	···	···	disk or low gravity?	? (too faint)
S Ori 71	disk	···	disk	disk
S Ori J053858.6−025228	no disk	disk	···	? (too faint)
S Ori J053949.5−023130	? (too faint)	···	disk	no disk
S Ori J053956.8−025315	? (too faint)	disk	···	? (too faint)

[a] Caballero et al. (2007). [b] Zapatero Osorio et al. (2007). [c] Scholz & Jayawardhana (2008). [d] Spectroscopically classified as a field dwarf rather than a cluster member (McGovern et al. 2004).

TABLE 4
Disk Fractions

M_J	Chamaeleon I	IC 348	σ Ori
0-2	$7/11 = 0.64^{+0.12}_{-0.15}$	$7/19 = 0.37^{+0.12}_{-0.09}$	$4/14 = 0.29^{+0.14}_{-0.09}$
2-4	$23/35 = 0.66^{+0.07}_{-0.09}$	$19/43 = 0.44^{+0.08}_{-0.07}$	$20/71 = 0.28^{+0.06}_{-0.05}$
4-6	$43/91 = 0.47^{+0.05}_{-0.05}$	$57/139 = 0.41^{+0.04}_{-0.04}$	$62/164 = 0.38^{+0.04}_{-0.04}$
6-8	$21/44 = 0.48^{+0.08}_{-0.07}$	$34/71 = 0.48^{+0.06}_{-0.06}$	$33/87 = 0.38^{+0.06}_{-0.05}$
8-10	$8/12 = 0.67^{+0.10}_{-0.15}$	$11/19 = 0.58^{+0.10}_{-0.11}$	$7/11 = 0.64^{+0.12}_{-0.15}$
10-12	$2/6 = 0.33^{+0.22}_{-0.12}$	···	$5/10 = 0.50^{+0.15}_{-0.14}$

Note. — The statistical errors in the disk fractions are computed in the manner described by Burgasser et al. (2003).



FIG. 1.— Fields in the σ Ori cluster that have been imaged with IRAC. In each of the four IRAC filters, the total exposure times were 80.4 s in the large fields ("shallow", Hernández et al. 2007a) and $\geq$ 1120 s in the smaller regions ("deep", Scholz & Jayawardhana 2008). The image is from DSS ($1^\circ \times 1^\circ$).



FIG. 2.— *Spitzer* IRAC color-color diagrams for the shallow and deep images of the σ Ori cluster (*left and right*). *Top*: Probable members of the cluster (*circles*) exhibit either neutral colors consistent with stellar photospheres or significantly redder colors that are indicative of circumstellar disks. *Bottom*: Among the other point sources in the IRAC data (*points*), candidate disk-bearing members of the cluster can be identified through their red colors ($[5.8] - [8.0] > 0.3$, $[3.6] - [4.5] > 0.15$, *dashed line*). Most of these candidates are probably galaxies based on their colors and magnitudes (see Fig. 3). Only objects with photometric errors less than 0.1 mag in all four bands are shown in these diagrams. The reddening vector is based on the extinction law from Flaherty et al. (2007).



FIG. 3.— *Spitzer* color-magnitude diagrams for the shallow and deep images of the σ Ori cluster (*left and right*). We show probable members of the cluster (*circles*) and sources from Figure 2 that have red colors ($[3.6] - [4.5] > 0.15$, $[4.5] - [5.8] > 0.15$, $[5.8] - [8.0] > 0.3$) and lack membership data (*points*). Most of the latter sources ere probably galaxies based on their faint magnitudes and very red colors. Only objects with photometric errors less than 0.1 mag in all four bands are shown in these diagrams. The 8 μm completeness limits are indicated (*dashed lines*). The reddening vector is based on the extinction law from Flaherty et al. (2007).



FIG. 4.— *Spitzer* color-magnitude diagrams for probable members of σ Ori in the shallow and deep images of the cluster (*top and bottom*). For comparison, we show a fit to the sequence of diskless members of Chamaeleon I after adjusting it to the distance of σ Ori (*left solid lines*). This fit is connected to the typical colors of T6 dwarfs for the magnitude of the T dwarf S Ori 70 (*dashed lines*). The 2σ scatter of the σ Ori sequence about this fit represents out adopted threshold for identifying sources that exhibit significant color excesses at brighter magnitudes ($[3.6] < 14.5$, *right solid lines*). At the fainter levels encompassing the brown dwarf candidates that were analyzed by Zapatero Osorio et al. (2007) and Scholz & Jayawardhana (2008), we have identified color excesses by considering the Poisson uncertainties in the colors (*error bars*), the size of the scatter in colors, and the general tendency to overestimate fluxes for sources with very low SNRs (§ 4.1). The brown dwarf candidates that we classify as "disk?" or "disk" in Table 3 are indicated (*triangles and circles*).



FIG. 5.— Disk fraction as a function of absolute J-band magnitude for probable members of the σ Ori cluster based on the IRAC color excesses in Figure 4. For comparison, we include the disk fractions measured from IRAC data in IC 348 and Chamaeleon I (Luhman et al. 2005, 2008; Lada et al. 2006). The typical magnitudes for spectral types of K5 ($\sim 1\ M_\odot$) and M6 ($\sim 0.1\ M_\odot$) are indicated.



FIG. 6.— Comparison of the IRAC magnitudes from Caballero et al. (2007) and Zapatero Osorio et al. (2007) to our measurements of the same objects. Both sets of data were measured from the shallow images of the σ Ori cluster. The errors reported by Caballero et al. (2007) and Zapatero Osorio et al. (2007) are indicated. The formal errors for our measurements range between 0.02–0.12 mag.



FIG. 7.— Comparison of the IRAC magnitudes from Scholz & Jayawardhana (2008) to our measurements of the same objects. Both sets of data were measured from the deep images of the σ Ori cluster. The errors reported by Scholz & Jayawardhana (2008) are indicated. The formal errors for our measurements are smaller by factors of 2–3 in most cases.



FIG. 8.— Comparison of the IRAC colors from Caballero et al. (2007), Zapatero Osorio et al. (2007), and Scholz & Jayawardhana (2008) to our measurements of the same objects from the same images considered in those studies, which are the shallow and deep images of the σ Ori cluster (*left and right*).



Fig. 9.— IRAC images at 5.8 and 8.0 μm (*odd and even columns*) centered on the positions of six of the candidate brown dwarfs in σ Ori that were discussed by Zapatero Osorio et al. (2007). Based on the shallow images, Zapatero Osorio et al. (2007) reported detections of excess emission that indicated the presence of disks for all of these sources. However, we find that the SNRs of these data are insufficient for reliably detecting excess emission. Deep images do show significant excesses for S Ori 56 and S Ori 60 and marginal excesses for S Ori 55 and S Ori 58 (Figure 4). Each image has a size of $30'' \times 30''$ and is displayed linearly from $F - 2\ \sigma$ to $F + 5\ \sigma$, where F and σ are the median and standard deviation of the background emission, respectively.



FIG. 10.— Deep IRAC images at 5.8 and 8.0 μm (*left and right*) centered on the positions of three of the candidate brown dwarfs in σ Ori that were discussed by Scholz & Jayawardhana (2008). Based on these images, Scholz & Jayawardhana (2008) reported detections of excess emission that indicated the presence of disks (their detection was only tentative for S Ori 70). However, we find that the SNRs of the 8 μm data are insufficient for reliably detecting excess emission. In the images at 5.8 μm, which have higher SNRs, S Ori 65 and S Ori J053949.5−023130 do not exhibit excesses. S Ori 70 does have an excess at 5.8 μm, but this measurement may be overestimated because of its low SNR (Beichman et al. 2003). Each image has a size of $30'' \times 30''$ and is displayed linearly from $F - 2\ \sigma$ to $F + 5\ \sigma$, where F and σ are the median and standard deviation of the background emission, respectively.